

**PARK7 Antibody**  
Rabbit mAb  
Catalog # AP90747

## Specification

---

### PARK7 Antibody - Product Information

Application	WB, IHC, FC, ICC, IP
Primary Accession	<a href="#">Q99497</a>
Reactivity	Rat
Clonality	Monoclonal
<b>Other Names</b>	
Protein DJ-1; SP22; Protein DJ-1; Oncogene DJ1; Parkinson disease protein 7; PARK7;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	19891 Da

### PARK7 Antibody - Additional Information

Dilution	WB~~1:1000 IHC~~1:100~500 FC~~1:10~50 ICC~~N/A IP~~N/A
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human PARK7
Description	Plays a role in regulating expression or stability of the mitochondrial uncoupling proteins SLC25A14 and SLC25A27 in dopaminergic neurons of the substantia nigra pars compacta and attenuates the oxidative stress induced by calcium entry into the neurons via L-type channels during pacemaking. It cooperates with Ras to increase cell transformation, it positively regulates transcription of the androgen receptor, and it may function as an indicator of oxidative stress.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

### PARK7 Antibody - Protein Information

Name PARK7 ([HGNC:16369](#))

## Function

Multifunctional protein with controversial molecular function which plays an important role in cell protection against oxidative stress and cell death acting as oxidative stress sensor and redox-sensitive chaperone and protease (PubMed: [12796482](http://www.uniprot.org/citations/12796482), PubMed: [17015834](http://www.uniprot.org/citations/17015834), PubMed: [18711745](http://www.uniprot.org/citations/18711745), PubMed: [19229105](http://www.uniprot.org/citations/19229105), PubMed: [20304780](http://www.uniprot.org/citations/20304780), PubMed: [25416785](http://www.uniprot.org/citations/25416785), PubMed: [26995087](http://www.uniprot.org/citations/26995087), PubMed: [28993701](http://www.uniprot.org/citations/28993701)). It is involved in neuroprotective mechanisms like the stabilization of NFE2L2 and PINK1 proteins, male fertility as a positive regulator of androgen signaling pathway as well as cell growth and transformation through, for instance, the modulation of NF-kappa-B signaling pathway (PubMed: [12612053](http://www.uniprot.org/citations/12612053), PubMed: [14749723](http://www.uniprot.org/citations/14749723), PubMed: [15502874](http://www.uniprot.org/citations/15502874), PubMed: [17015834](http://www.uniprot.org/citations/17015834), PubMed: [18711745](http://www.uniprot.org/citations/18711745), PubMed: [21097510](http://www.uniprot.org/citations/21097510), PubMed: [21097510](http://www.uniprot.org/citations/21097510)). Has been described as a protein and nucleotide deglycase that catalyzes the deglycation of the Maillard adducts formed between amino groups of proteins or nucleotides and reactive carbonyl groups of glyoxals (PubMed: [25416785](http://www.uniprot.org/citations/25416785), PubMed: [28596309](http://www.uniprot.org/citations/28596309)). But this function is rebutted by other works (PubMed: [27903648](http://www.uniprot.org/citations/27903648), PubMed: [31653696](http://www.uniprot.org/citations/31653696)). As a protein deglycase, repairs methylglyoxal- and glyoxal-glycated proteins, and releases repaired proteins and lactate or glycolate, respectively. Deglycates cysteine, arginine and lysine residues in proteins, and thus reactivates these proteins by reversing glycation by glyoxals. Acts on early glycation intermediates (hemithioacetals and aminocarbonyls), preventing the formation of advanced glycation endproducts (AGE) that cause irreversible damage (PubMed: [25416785](http://www.uniprot.org/citations/25416785), PubMed: [26995087](http://www.uniprot.org/citations/26995087), PubMed: [28013050](http://www.uniprot.org/citations/28013050), PubMed: [28013050](http://www.uniprot.org/citations/28013050)). Also functions as a nucleotide deglycase able to repair glycated guanine in the free nucleotide pool (GTP, GDP, GMP, dGTP) and in DNA and RNA. Is thus involved in a major nucleotide repair system named guanine glycation repair (GG repair), dedicated to reversing methylglyoxal and glyoxal damage via nucleotide sanitization and direct nucleic acid repair (PubMed: [28596309](http://www.uniprot.org/citations/28596309), PubMed: [28596309](http://www.uniprot.org/citations/28596309)). Protects histones from adduction by methylglyoxal, controls the levels of methylglyoxal- derived argininine modifications on chromatin (PubMed: [30150385](http://www.uniprot.org/citations/30150385), PubMed: [30150385](http://www.uniprot.org/citations/30150385)). Able to remove the glycations and restore histone 3, histone glycation disrupts both local and global chromatin architecture by altering histone-DNA interactions as well as histone acetylation and ubiquitination levels (PubMed: [30150385](http://www.uniprot.org/citations/30150385), PubMed: [30894531](http://www.uniprot.org/citations/30894531), PubMed: [30894531](http://www.uniprot.org/citations/30894531)). Displays a very low glyoxalase activity that may reflect its deglycase activity (PubMed: [22523093](http://www.uniprot.org/citations/22523093), PubMed: [22523093](http://www.uniprot.org/citations/22523093), PubMed: [28993701](http://www.uniprot.org/citations/28993701), PubMed: [28993701](http://www.uniprot.org/citations/28993701), PubMed: [31653696](http://www.uniprot.org/citations/31653696), PubMed: [31653696](http://www.uniprot.org/citations/31653696)). Eliminates hydrogen peroxide and protects cells against hydrogen peroxide-induced cell death (PubMed: [16390825](http://www.uniprot.org/citations/16390825), PubMed: [16390825](http://www.uniprot.org/citations/16390825)). Required for correct mitochondrial morphology and function as well as for autophagy of dysfunctional mitochondria (PubMed: [16632486](http://www.uniprot.org/citations/16632486), PubMed: [16632486](http://www.uniprot.org/citations/16632486), PubMed: [19229105](http://www.uniprot.org/citations/19229105), PubMed: [19229105](http://www.uniprot.org/citations/19229105)).

target="\_blank">19229105</a>). Plays a role in regulating expression or stability of the mitochondrial uncoupling proteins SLC25A14 and SLC25A27 in dopaminergic neurons of the substantia nigra pars compacta and attenuates the oxidative stress induced by calcium entry into the neurons via L-type channels during pacemaking (PubMed:<a href="http://www.uniprot.org/citations/18711745" target="\_blank">18711745</a>). Regulates astrocyte inflammatory responses, may modulate lipid raft-dependent endocytosis in astrocytes and neuronal cells (PubMed:<a href="http://www.uniprot.org/citations/23847046" target="\_blank">23847046</a>). In pancreatic islets, involved in the maintenance of mitochondrial reactive oxygen species (ROS) levels and glucose homeostasis in an age- and diet dependent manner. Protects pancreatic beta cells from cell death induced by inflammatory and cytotoxic setting (By similarity). Binds to a number of mRNAs containing multiple copies of GG or CC motifs and partially inhibits their translation but dissociates following oxidative stress (PubMed:<a href="http://www.uniprot.org/citations/18626009" target="\_blank">18626009</a>). Metal-binding protein able to bind copper as well as toxic mercury ions, enhances the cell protection mechanism against induced metal toxicity (PubMed:<a href="http://www.uniprot.org/citations/23792957" target="\_blank">23792957</a>). In macrophages, interacts with the NADPH oxidase subunit NCF1 to direct NADPH oxidase-dependent ROS production, and protects against sepsis (By similarity).

### Cellular Location

Cell membrane {ECO:0000250|UniProtKB:Q99LX0}; Lipid-anchor {ECO:0000250|UniProtKB:Q99LX0}. Cytoplasm. Nucleus. Membrane raft {ECO:0000250|UniProtKB:O88767}. Mitochondrion. Endoplasmic reticulum. Note=Under normal conditions, located predominantly in the cytoplasm and, to a lesser extent, in the nucleus and mitochondrion. Translocates to the mitochondrion and subsequently to the nucleus in response to oxidative stress and exerts an increased cytoprotective effect against oxidative damage (PubMed:18711745). Detected in tau inclusions in brains from neurodegenerative disease patients (PubMed:14705119). Membrane raft localization in astrocytes and neuronal cells requires palmitoylation

### Tissue Location


Highly expressed in pancreas, kidney, skeletal muscle, liver, testis and heart. Detected at slightly lower levels in placenta and brain (at protein level). Detected in astrocytes, Sertoli cells, spermatogonia, spermatids and spermatozoa. Expressed by pancreatic islets at higher levels than surrounding exocrine tissues (PubMed:22611253).


## PARK7 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## PARK7 Antibody - Images





Western blot analysis of PARK7 expression in HeLa cell lysate.