

HDAC7 Antibody
Rabbit mAb
Catalog # AP90901**Specification**

HDAC7 Antibody - Product Information

Application	WB, FC
Primary Accession	Q8WUI4
Reactivity	Rat
Clonality	Monoclonal
Other Names	
HD 7a; HD7; HDAC 7; HDAC 7A; Hdac7; HDAC7A; Histone deacetylase 7; Histone deacetylase 7A;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	102927 Da

HDAC7 Antibody - Additional Information

Dilution	WB~~1:1000 FC~~1:10~50
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human HDAC7
Description	Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes. Involved in muscle maturation by repressing transcription of myocyte enhancer factors such as MEF2A, MEF2B and MEF2C.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

HDAC7 Antibody - Protein Information**Name** HDAC7**Synonyms** HDAC7A**Function**

Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4) (By similarity). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events (By similarity). Histone deacetylases act via the formation of large multiprotein complexes (By similarity). Involved in muscle maturation by repressing transcription of myocyte enhancer factors such as MEF2A, MEF2B and MEF2C (By similarity). During muscle differentiation, it shuttles into the cytoplasm, allowing the expression of myocyte enhancer factors (By similarity). May be involved in Epstein-Barr virus (EBV) latency, possibly by repressing the viral BZLF1 gene (PubMed: 12239305). Positively regulates the transcriptional repressor activity of FOXP3 (PubMed: 17360565). Serves as a corepressor of RARA, causing its deacetylation and inhibition of RARE DNA element binding (PubMed: 28167758). In association with RARA, plays a role in the repression of microRNA-10a and thereby in the inflammatory response (PubMed: 28167758). Also acetylates non-histone proteins, such as ALKBH5 (PubMed: 37369679).

Cellular Location

Nucleus. Cytoplasm Note=In the nucleus, it associates with distinct subnuclear dot-like structures (PubMed:11262386). Shuttles between the nucleus and the cytoplasm (PubMed:16980613). In muscle cells, it shuttles into the cytoplasm during myocyte differentiation (By similarity). The export to cytoplasm depends on the interaction with the 14-3-3 protein YWHAE and is due to its phosphorylation (PubMed:16980613) {ECO:0000250|UniProtKB:Q8C2B3, ECO:0000269|PubMed:11262386, ECO:0000269|PubMed:16980613}


HDAC7 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HDAC7 Antibody - Images





Western blot analysis of HDAC7 expression in HeLa cell lysate.