

# ADRBK2 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP9092a

## **Specification**

## ADRBK2 Antibody (N-term) - Product Information

Application WB, IHC-P,E **Primary Accession** P35626 Reactivity Human Host Rabbit Clonality **Polyclonal** Isotype Rabbit IgG Calculated MW 79710 **Antigen Region** 29-57

### ADRBK2 Antibody (N-term) - Additional Information

#### Gene ID 157

#### **Other Names**

Beta-adrenergic receptor kinase 2, Beta-ARK-2, G-protein-coupled receptor kinase 3, ADRBK2, BARK2, GRK3

#### Target/Specificity

This ADRBK2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 29-57 amino acids from the N-terminal region of human ADRBK2.

## **Dilution**

WB~~1:1000 IHC-P~~1:50~100

### **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

### **Precautions**

ADRBK2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

### ADRBK2 Antibody (N-term) - Protein Information

## Name GRK3 (HGNC:290)

Function Specifically phosphorylates the agonist-occupied form of the beta-adrenergic and



closely related receptors.

#### **Cellular Location**


Postsynapse {ECO:0000250|UniProtKB:P26819}. Presynapse {ECO:0000250|UniProtKB:P26819}

### ADRBK2 Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

## ADRBK2 Antibody (N-term) - Images



Western blot analysis of ADRBK2 (arrow) using rabbit polyclonal ADRBK2 Antibody (N-term) (Cat. #AP9092a). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the ADRBK2 gene (Lane 2) .



Formalin-fixed and paraffin-embedded human brain tissue reacted with ADRBK2 Antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining.



Tel: 858.875.1900 Fax: 858.875.1999

This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

### ADRBK2 Antibody (N-term) - Background

ADRBK2 is the beta-adrenergic receptor kinase specifically phosphorylates the agonist-occupied form of the beta-adrenergic and related G protein-coupled receptors. Overall, the beta adrenergic receptor kinase 2 has 85% amino acid similarity with beta adrenergic receptor kinase 1, with the protein kinase catalytic domain having 95% similarity. These data suggest the existence of a family of receptor kinases which may serve broadly to regulate receptor function.

## ADRBK2 Antibody (N-term) - References

Gratacos, M., et.al., Am. J. Med. Genet. B Neuropsychiatr. Genet. 150B (6), 808-816 (2009) Rao, J.S., et.al., Int. J. Neuropsychopharmacol. 12 (6), 851-860 (2009)