

**TSG101 Antibody**  
**Rabbit mAb**  
**Catalog # AP90984****Specification**

---

**TSG101 Antibody - Product Information**

Application	WB, IHC, FC, ICC
Primary Accession	<a href="#">Q99816</a>
Reactivity	Rat
Clonality	Monoclonal

**Other Names**

TSG101; ESCRT-I complex subunit TSG101; Tumor susceptibility gene 10; Tumor susceptibility protein; VPS23; TSG10;

Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	43944 Da

**TSG101 Antibody - Additional Information**

Dilution	WB~~1:1000 IHC~~1:100~500 FC~~1:10~50 ICC~~N/A
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human TSG101
Description	Component of the ESCRT-I complex, a regulator of vesicular trafficking process. Binds to ubiquitinated cargo proteins and is required for the sorting of endocytic ubiquitinated cargos into multivesicular bodies (MVBs). Mediates the association between the ESCRT-0 and ESCRT-I complex. Required for completion of cytokinesis; the function requires CEP55.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

**TSG101 Antibody - Protein Information**

**Name** TSG101

**Function**

Component of the ESCRT-I complex, a regulator of vesicular trafficking process. Binds to ubiquitinated cargo proteins and is required for the sorting of endocytic ubiquitinated cargos into

multivesicular bodies (MVBs). Mediates the association between the ESCRT-0 and ESCRT-I complex. Required for completion of cytokinesis; the function requires CEP55. May be involved in cell growth and differentiation. Acts as a negative growth regulator. Involved in the budding of many viruses through an interaction with viral proteins that contain a late-budding motif P-[ST]-A-P. This interaction is essential for viral particle budding of numerous retroviruses. Required for the exosomal release of SDCBP, CD63 and syndecan (PubMed:<a href="http://www.uniprot.org/citations/22660413" target="\_blank">22660413</a>). It may also play a role in the extracellular release of microvesicles that differ from the exosomes (PubMed:<a href="http://www.uniprot.org/citations/22315426" target="\_blank">22315426</a>).

#### Cellular Location

Cytoplasm. Early endosome membrane; Peripheral membrane protein; Cytoplasmic side. Late endosome membrane; Peripheral membrane protein. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Midbody, Midbody ring. Nucleus. Note=Mainly cytoplasmic. Membrane- associated when active and soluble when inactive. Nuclear localization is cell cycle-dependent. Interaction with CEP55 is required for localization to the midbody during cytokinesis

#### Tissue Location


Heart, brain, placenta, lung, liver, skeletal, kidney and pancreas

#### TSG101 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### TSG101 Antibody - Images



Western blot analysis of TSG101 expression in Jurkat cell lysate.