

## **iNOS Antibody**

Rabbit mAb Catalog # AP90991

### **Specification**

### **iNOS Antibody - Product Information**

Application WB, ICC, IP
Primary Accession P35228
Clonality Monoclonal

**Other Names** 

HEP-NOS; Hepatocyte NOS; Inducible NO synthase; Inducible NOS; iNOS; NOS2;

Isotype Rabbit IgG
Host Rabbit
Calculated MW 131117 Da

### **iNOS Antibody - Additional Information**

Dilution **WB~~1:1000** 

ICC~~N/A IP~~N/A

Purification Affinity-chromatography

Immunogen A synthesized peptide derived from human

**iNOS** 

Description iNOS Produces nitric oxide (NO) which is a

messenger molecule with diverse functions throughout the body. In macrophages, NO mediates tumoricidal and bactericidal

actions.

Storage Condition and Buffer Rabbit IgG in phosphate buffered saline,

pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid

freeze / thaw cycle.

## **iNOS Antibody - Protein Information**

Name NOS2 (HGNC:7873)

Synonyms NOS2A

#### **Function**

Produces nitric oxide (NO) which is a messenger molecule with diverse functions throughout the body (PubMed:<a href="http://www.uniprot.org/citations/7504305"

target="\_blank">7504305</a>, PubMed:<a href="http://www.uniprot.org/citations/7531687" target="\_blank">7531687</a>, PubMed:<a href="http://www.uniprot.org/citations/7544004" target="\_blank">7544004</a>, PubMed:<a href="http://www.uniprot.org/citations/7682706" target="\_blank">7682706</a>). In macrophages, NO mediates tumoricidal and bactericidal actions. Also has nitrosylase activity and mediates cysteine S-nitrosylation of cytoplasmic target



Tel: 858.875.1900 Fax: 858.875.1999

proteins such PTGS2/COX2 (By similarity). As component of the iNOS-S100A8/9 transnitrosylase complex involved in the selective inflammatory stimulus-dependent S-nitrosylation of GAPDH on 'Cys-247' implicated in regulation of the GAIT complex activity and probably multiple targets including ANXA5, EZR, MSN and VIM (PubMed:<a

href="http://www.uniprot.org/citations/25417112" target="\_blank">25417112</a>). Involved in inflammation, enhances the synthesis of pro-inflammatory mediators such as IL6 and IL8 (PubMed:<a href="http://www.uniprot.org/citations/19688109" target="\_blank">19688109</a>).

## **Cellular Location**

Cytoplasm, cytosol. Note=Localizes as discrete foci scattered throughout the cytosol and in the presence of SPSB1 and SPSB4, exhibits a more diffuse cytosolic localization.

#### **Tissue Location**


Expressed in the liver, retina, bone cells and airway epithelial cells of the lung. Not expressed in the platelets Expressed in chondrocytes (PubMed:7504305)

# **iNOS Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

## iNOS Antibody - Images



Western blot analysis of iNOS expression in Human fetal brain lysate.