

Interferon beta Antibody

Rabbit mAb Catalog # AP91121

Specification

Interferon beta Antibody - Product Information

Application WB, IP
Primary Accession P01574
Clonality Monoclonal

Other Names

Interferon beta; Interferon β; IFN-beta; IFN-β; IFNβ; Fibroblast interferon; IFB; IFNB;

Isotype Rabbit IgG
Host Rabbit
Calculated MW 22294 Da

Interferon beta Antibody - Additional Information

Dilution WB~~1:1000

IP~~N/A

Purification Affinity-chromatography

Immunogen A synthesized peptide derived from human

Interferon beta

Description The type I interferon (IFN) family includes

IFN-β1 and IFN-α1 through IFN-α13 in humans and IFN-α1 through IFN-α14 in mice. Type I IFN is produced following detection of pathogen-associated molecular patterns (PAMPs) and is

important for induction of antiviral genes, activation of dendritic cells, and initiation of adaptive immunity. Has antiviral,

antibacterial and anticancer activities. Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide

and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid

freeze / thaw cycle.

Interferon beta Antibody - Protein Information

Name IFNB1 (HGNC:5434)

Storage Condition and Buffer

Synonyms IFB, IFNB

Function

Type I interferon cytokine that plays a key role in the innate immune response to infection, developing tumors and other inflammatory stimuli (PubMed:10049744, PubMed:<a



href="http://www.uniprot.org/citations/10556041" target=" blank">10556041, PubMed:6157094, PubMed:6171735, PubMed:7665574, PubMed:8027027, PubMed:8969169). Signals via binding to high-affinity (IFNAR2) and low-affinity (IFNAR1) heterodimeric receptor, activating the canonical Jak-STAT signaling pathway resulting in transcriptional activation or repression of interferon-regulated genes that encode the effectors of the interferon response, such as antiviral proteins, regulators of cell proliferation and differentiation, and immunoregulatory proteins (PubMed:10049744, PubMed: 10556041, PubMed: 7665574, PubMed: 8027027, PubMed:8969169). Signals mostly via binding to a IFNAR1-IFNAR2 heterodimeric receptor, but can also function with IFNAR1 alone and independently of Jak-STAT pathways (By similarity). Elicits a wide variety of responses, including antiviral and antibacterial activities, and can regulate the development of B-cells, myelopoiesis and lipopolysaccharide (LPS)- inducible production of tumor necrosis factor (By similarity). Plays a role in neuronal homeostasis by regulating dopamine turnover and protecting dopaminergic neurons: acts by promoting neuronal autophagy and alpha-synuclein clearance, thereby preventing dopaminergic neuron loss (By similarity). IFNB1 is more potent than interferon-alpha (IFN- alpha) in inducing the apoptotic and antiproliferative pathways required for control of tumor cell growth (By similarity).

Cellular Location Secreted.

Interferon beta Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Interferon beta Antibody - Images



