

ADAM17 Antibody
Rabbit mAb
Catalog # AP91129**Specification**

ADAM17 Antibody - Product Information

| | |
|-----------------------------|------------------------|
| Application | WB, FC, ICC, IP |
| Primary Accession | P78536 |
| Reactivity | Rat |
| Clonality | Monoclonal |
| Other Names | |
| CD156b; ADAM17; CSVP; TACE; | |
| Isotype | Rabbit IgG |
| Host | Rabbit |
| Calculated MW | 93021 Da |

ADAM17 Antibody - Additional Information

| | |
|------------------------------|--|
| Dilution | WB~~1:1000 FC~~1:10~50 ICC~~N/A IP~~N/A |
| Purification | Affinity-chromatography |
| Immunogen | A synthesized peptide derived from human ADAM17 |
| Description | TACE is responsible for the shedding of EGFR ligands such as amphiregulin and TNF- α . Some tumors have hyperactivated EGFR due to upregulated TNF- α production and upregulated TACE, making TACE a potential target for drug development. TACE activates Notch in a ligand-independent manner and has been shown to play a role in the development of the Drosophila nervous system. |
| Storage Condition and Buffer | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle. |

ADAM17 Antibody - Protein Information**Name** ADAM17 ([HGNC:195](#))**Synonyms** CSVP, TACE**Function**

Transmembrane metalloprotease which mediates the ectodomain shedding of a myriad of transmembrane proteins including adhesion proteins, growth factor precursors and cytokines important for inflammation and immunity (PubMed:24226769, PubMed:24227843, PubMed:28060820, PubMed:28923481). Cleaves the membrane-bound precursor of TNF-alpha to its mature soluble form (PubMed:36078095, PubMed:9034191). Responsible for the proteolytical release of soluble JAM3 from endothelial cells surface (PubMed:20592283). Responsible for the proteolytic release of several other cell-surface proteins, including p75 TNF-receptor, interleukin 1 receptor type II, p55 TNF- receptor, transforming growth factor-alpha, L-selectin, growth hormone receptor, MUC1 and the amyloid precursor protein (PubMed:12441351). Acts as an activator of Notch pathway by mediating cleavage of Notch, generating the membrane-associated intermediate fragment called Notch extracellular truncation (NEXT) (PubMed:24226769). Plays a role in the proteolytic processing of ACE2 (PubMed:24227843). Plays a role in hemostasis through shedding of GP1BA, the platelet glycoprotein Ib alpha chain (By similarity). Mediates the proteolytic cleavage of LAG3, leading to release the secreted form of LAG3 (By similarity). Mediates the proteolytic cleavage of IL6R, leading to the release of secreted form of IL6R (PubMed:26876177, PubMed:28060820). Mediates the proteolytic cleavage and shedding of FCGR3A upon NK cell stimulation, a mechanism that allows for increased NK cell motility and detachment from opsonized target cells. Cleaves TREM2, resulting in shedding of the TREM2 ectodomain (PubMed:28923481).

Cellular Location

Cell membrane; Single-pass type I membrane protein

Tissue Location

Ubiquitously expressed. Expressed at highest levels in adult heart, placenta, skeletal muscle, pancreas, spleen, thymus, prostate, testes, ovary and small intestine, and in fetal brain, lung, liver and kidney. Expressed in natural killer cells (at protein level) (PubMed:24337742).


ADAM17 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ADAM17 Antibody - Images





Western blot analysis of ADAM17 expression in Hela cell lysate.