

SDHA Antibody

Rabbit mAb Catalog # AP91182

Specification

SDHA Antibody - Product Information

Application Primary Accession Reactivity Clonality Other Names PGL5; SDH1; SDH2; SDHA; SDHF;	WB, IHC, FC, ICC, IP <u>P31040</u> Rat Monoclonal
lsotype Host Calculated MW	Rabbit IgG Rabbit 72692 Da
SDHA Antibody - Additional Information	
Dilution	WB~~1:1000 IHC~~1:100~500 FC~~1:10~50 ICC~~N/A IP~~N/A
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human SDHA
Description	Flavoprotein (FP) subunit of succinate dehydrogenase (SDH) that is involved in complex II of the mitochondrial electron transport chain and is responsible for transferring electrons from succinate to

Storage Condition and Buffer

SDHA Antibody - Protein Information

Name SDHA

Synonyms SDH2, SDHF

Function

Flavoprotein (FP) subunit of succinate dehydrogenase (SDH) that is involved in complex II of the mitochondrial electron transport chain and is responsible for transferring electrons from succinate to ubiquinone (coenzyme Q) (PubMed:http://www.uniprot.org/citations/10746566

ubiquinone.

freeze / thaw cycle.

Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid



target="_blank">10746566, PubMed:24781757). SDH also oxidizes malate to the non-canonical enol form of oxaloacetate, enol- oxaloacetate (By similarity). Enol-oxaloacetate, which is a potent inhibitor of the succinate dehydrogenase activity, is further isomerized into keto-oxaloacetate (By similarity). Can act as a tumor suppressor (PubMed:20484225).

Cellular Location


Mitochondrion inner membrane; Peripheral membrane protein; Matrix side

SDHA Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

SDHA Antibody - Images



Western blot analysis of SDHA expression in (1) HeLa cell lysate; (2) Mouse brain lysate; (3) Rat brain lysate;