

**Apg3 (Atg3) Antibody**  
**Rabbit mAb**  
**Catalog # AP91265****Specification**

---

**Apg3 (Atg3) Antibody - Product Information**

Application	WB, IHC, ICC
Primary Accession	<a href="#">Q9NT62</a>
Reactivity	Rat
Clonality	Monoclonal

**Other Names**

APG3 autophagy 3 like; APG3 like; APG3, *S. cerevisiae*, homolog of; APG3-like; APG3L; Apg3p; ATG3 autophagy related 3 homolog;

Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	35864 Da

**Apg3 (Atg3) Antibody - Additional Information**

Dilution	WB~~1:1000 IHC~~1:100~500 ICC~~N/A
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Apg3 (Atg3)
Description	E2-like enzyme involved in autophagy and mitochondrial homeostasis. Catalyzes the conjugation of ATG8-like proteins (GABARAP, GABARAPL1, GABARAPL2 or MAP1LC3A) to phosphatidylethanolamine (PE). Catalyzing the conjugation of ATG12 to itself, ATG12 conjugation to ATG3 playing a role in mitochondrial homeostasis but not in autophagy. ATG7 (E1-like enzyme) facilitates this reaction by forming an E1-E2 complex with ATG3.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

**Apg3 (Atg3) Antibody - Protein Information**

**Name** ATG3 ([HGNC:20962](#))

**Synonyms** APG3, APG3L

**Function**

E2 conjugating enzyme that catalyzes the covalent conjugation of the C-terminal Gly of ATG8-like proteins (GABARAP, GABARAPL1, GABARAPL2 or MAP1LC3A) to the amino group of phosphatidylethanolamine (PE)-containing lipids in the membrane resulting in membrane-bound ATG8-like proteins which is one of the key steps in the development of autophagic isolation membranes during autophagosome formation (PubMed:<a href="http://www.uniprot.org/citations/24191030" target="\_blank">24191030</a>, PubMed:<a href="http://www.uniprot.org/citations/33446636" target="\_blank">33446636</a>, PubMed:<a href="http://www.uniprot.org/citations/37252361" target="\_blank">37252361</a>). Cycles back and forth between binding to ATG7 for loading with the ATG8-like proteins and binding to E3 enzyme, composed of ATG12, ATG5 and ATG16L1 to promote ATG8-like proteins lipidation (PubMed:<a href="http://www.uniprot.org/citations/11825910" target="\_blank">11825910</a>, PubMed:<a href="http://www.uniprot.org/citations/12207896" target="\_blank">12207896</a>, PubMed:<a href="http://www.uniprot.org/citations/12890687" target="\_blank">12890687</a>, PubMed:<a href="http://www.uniprot.org/citations/16704426" target="\_blank">16704426</a>, PubMed:<a href="http://www.uniprot.org/citations/24186333" target="\_blank">24186333</a>). Also plays a role as a membrane curvature sensor that facilitates LC3/GABARAP lipidation by sensing local membrane stress associated with lipid-packing defects as occurs with high molar proportions of conical lipids or strident membrane curvature (By similarity). Interacts with negatively-charged membranes promoting membrane tethering and enhancing LC3/GABARAP lipidation (PubMed:<a href="http://www.uniprot.org/citations/29142222" target="\_blank">29142222</a>). Also acts as an autocatalytic E2-like enzyme by catalyzing the conjugation of ATG12 to itself in an ATG7-dependent manner, this complex thus formed, plays a role in mitochondrial homeostasis but not in autophagy (By similarity). ATG12- ATG3 conjugation promotes late endosome to lysosome trafficking and basal autophagosome maturation via its interaction with PDCD6IP (By similarity). ATG12-ATG3 conjugate is also formed upon vaccinia virus infection, leading to the disruption the cellular autophagy which is not necessary for vaccinia survival and proliferation (By similarity). Promotes primary ciliogenesis by removing OFD1 from centriolar satellites via the autophagic pathway (By similarity).

**Cellular Location**

Cytoplasm.


**Tissue Location**


Widely expressed, with a highest expression in heart, skeletal muscle, kidney, liver and placenta

**Apg3 (Atg3) Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Apg3 (Atg3) Antibody - Images**



Western blot analysis of Apg3 (Atg3) expression in Jurkat cell lysate.