

GATA3 Antibody

Rabbit mAb Catalog # AP91270

Specification

GATA3 Antibody - Product Information

Application WB, IHC, FC, ICC, CHIP

Primary Accession
Clonality
Monoclonal
Other Names

GATA-binding factor 3; GATA3;

Isotype Rabbit IgG
Host Rabbit
Calculated MW 47916 Da

GATA3 Antibody - Additional Information

Dilution WB~~1:1000

IHC~~1:100~500 FC~~1:10~50 ICC~~N/A CHIP~~N/A

Purification Affinity-chromatography

Immunogen A synthesized peptide derived from human

GATA3

Description Transcriptional activator which binds to

the enhancer of the T-cell receptor alpha and delta genes. Binds to the consensus

sequence 5'-AGATAG-3'.

Storage Condition and Buffer Rabbit IgG in phosphate buffered saline,

pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid

freeze / thaw cycle.

GATA3 Antibody - Protein Information

Name GATA3

Function

Transcriptional activator which binds to the enhancer of the T-cell receptor alpha and delta genes. Binds to the consensus sequence 5'-AGATAG-3'. Required for the T-helper 2 (Th2) differentiation process following immune and inflammatory responses. Positively regulates ASB2 expression (By similarity). Coordinates macrophage transcriptional activation and UCP2-dependent metabolic reprogramming in response to IL33. Upon tissue injury, acts downstream of IL33 signaling to drive differentiation of inflammation-resolving alternatively activated macrophages.

Cellular Location



Nucleus.

Tissue Location


T-cells and endothelial cells.

GATA3 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

GATA3 Antibody - Images



Western blot analysis of GATA3 expression in (1) SH-SY5Y cell lysate; (2) Jurkat cell lysate.