

Drosha Antibody
Rabbit mAb
Catalog # AP91300

Specification

Drosha Antibody - Product Information

Application	WB, IHC, FC, ICC
Primary Accession	Q9NRR4
Clonality	Monoclonal
Other Names	
Ribonuclease 3 (EC:3.1.26.3); Protein Drosha; Ribonuclease III; RNase III; p241; DROSHA; RN3; RNASE3L; RNASEN;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	159316 Da

Drosha Antibody - Additional Information

Dilution	WB~~1:1000 IHC~~1:100~500 FC~~1:10~50 ICC~~N/A
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Drosha
Description	Drosha was identified as a nuclear RNase III that catalyzes the initial step of microRNA (miRNA) processing. This enzyme processes the long primary transcript pri-miRNAs into stem-looped pre-miRNAs. Interference of Drosha results in the increase of pri-miRNAs and the decrease of pre-miRNAs. Drosha exists in a multiprotein complex called Microprocessor along with other components such as DGCR8. Drosha, along with DGCR8, is necessary for miRNA biogenesis.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Drosha Antibody - Protein Information

Name DROSHA

Synonyms RN3, RNASE3L, RNASEN

Function

Ribonuclease III double-stranded (ds) RNA-specific endoribonuclease that is involved in the initial step of microRNA (miRNA) biogenesis. Component of the microprocessor complex that is required to process primary miRNA transcripts (pri-miRNAs) to release precursor miRNA (pre-miRNA) in the nucleus. Within the microprocessor complex, DROSHA cleaves the 3' and 5' strands of a stem-loop in pri- miRNAs (processing center 11 bp from the dsRNA-ssRNA junction) to release hairpin-shaped pre-miRNAs that are subsequently cut by the cytoplasmic DICER to generate mature miRNAs. Involved also in pre-rRNA processing. Cleaves double-strand RNA and does not cleave single-strand RNA. Involved in the formation of GW bodies. Plays a role in growth homeostasis in response to autophagy in motor neurons (By similarity).

Cellular Location

Nucleus. Nucleus, nucleolus. Cytoplasm {ECO:0000250|UniProtKB:Q5HZJ0}. Note=A fraction is translocated to the nucleolus during the S phase of the cell cycle. Localized in GW bodies (GWBs), also known as P-bodies.

Tissue Location


Ubiquitous..

Drosha Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Drosha Antibody - Images



Western blot analysis of Drosha expression in 293 cell lysate.