

**ITCH Antibody**  
**Rabbit mAb**  
**Catalog # AP91387****Specification****ITCH Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">Q96J02</a>
Reactivity	Rat
Clonality	Monoclonal
<b>Other Names</b>	ADMFD; AIF4; AIP4; Atrophin 1 interacting protein 4; Itchy homolog E3 ubiquitin protein ligase; NAPP1; NFE2 associated polypeptide 1; Ubiquitin protein ligase ITCH;
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	102803 Da

**ITCH Antibody - Additional Information**

Dilution	WB~~1:1000
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human ITCH
Description	Involved in the negative regulation of MAVS-dependent cellular antiviral responses. Ubiquitinates MAVS through 'Lys-48'-linked conjugation resulting in MAVS proteasomal degradation. Involved in the regulation of apoptosis and reactive oxygen species levels through the ubiquitination and proteasomal degradation of TXNIP. Mediates the antiapoptotic activity of epidermal growth factor through the ubiquitination and proteasomal degradation of p15 BID.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

**ITCH Antibody - Protein Information****Name** ITCH**Function**

Acts as an E3 ubiquitin-protein ligase which accepts ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and then directly transfers the ubiquitin to

targeted substrates (PubMed:<a href="http://www.uniprot.org/citations/11046148" target="\_blank">11046148</a>, PubMed:<a href="http://www.uniprot.org/citations/14602072" target="\_blank">14602072</a>, PubMed:<a href="http://www.uniprot.org/citations/15051726" target="\_blank">15051726</a>, PubMed:<a href="http://www.uniprot.org/citations/16387660" target="\_blank">16387660</a>, PubMed:<a href="http://www.uniprot.org/citations/17028573" target="\_blank">17028573</a>, PubMed:<a href="http://www.uniprot.org/citations/18718448" target="\_blank">18718448</a>, PubMed:<a href="http://www.uniprot.org/citations/18718449" target="\_blank">18718449</a>, PubMed:<a href="http://www.uniprot.org/citations/19116316" target="\_blank">19116316</a>, PubMed:<a href="http://www.uniprot.org/citations/19592251" target="\_blank">19592251</a>, PubMed:<a href="http://www.uniprot.org/citations/19881509" target="\_blank">19881509</a>, PubMed:<a href="http://www.uniprot.org/citations/20068034" target="\_blank">20068034</a>, PubMed:<a href="http://www.uniprot.org/citations/20392206" target="\_blank">20392206</a>, PubMed:<a href="http://www.uniprot.org/citations/20491914" target="\_blank">20491914</a>, PubMed:<a href="http://www.uniprot.org/citations/23146885" target="\_blank">23146885</a>, PubMed:<a href="http://www.uniprot.org/citations/24790097" target="\_blank">24790097</a>, PubMed:<a href="http://www.uniprot.org/citations/25631046" target="\_blank">25631046</a>). Catalyzes 'Lys-29'-, 'Lys-48'- and 'Lys-63'-linked ubiquitin conjugation (PubMed:<a href="http://www.uniprot.org/citations/17028573" target="\_blank">17028573</a>, PubMed:<a href="http://www.uniprot.org/citations/18718448" target="\_blank">18718448</a>, PubMed:<a href="http://www.uniprot.org/citations/19131965" target="\_blank">19131965</a>, PubMed:<a href="http://www.uniprot.org/citations/19881509" target="\_blank">19881509</a>). Involved in the control of inflammatory signaling pathways (PubMed:<a href="http://www.uniprot.org/citations/19131965" target="\_blank">19131965</a>). Essential component of a ubiquitin-editing protein complex, comprising also TNFAIP3, TAX1BP1 and RNF11, that ensures the transient nature of inflammatory signaling pathways (PubMed:<a href="http://www.uniprot.org/citations/19131965" target="\_blank">19131965</a>). Promotes the association of the complex after TNF stimulation (PubMed:<a href="http://www.uniprot.org/citations/19131965" target="\_blank">19131965</a>). Once the complex is formed, TNFAIP3 deubiquitinates 'Lys-63' polyubiquitin chains on RIPK1 and catalyzes the formation of 'Lys-48'-polyubiquitin chains (PubMed:<a href="http://www.uniprot.org/citations/19131965" target="\_blank">19131965</a>). This leads to RIPK1 proteasomal degradation and consequently termination of the TNF- or LPS-mediated activation of NFkB1 (PubMed:<a href="http://www.uniprot.org/citations/19131965" target="\_blank">19131965</a>). Ubiquitinates RIPK2 by 'Lys-63'-linked conjugation and influences NOD2-dependent signal transduction pathways (PubMed:<a href="http://www.uniprot.org/citations/19592251" target="\_blank">19592251</a>). Regulates the transcriptional activity of several transcription factors, and probably plays an important role in the regulation of immune response (PubMed:<a href="http://www.uniprot.org/citations/18718448" target="\_blank">18718448</a>, PubMed:<a href="http://www.uniprot.org/citations/20491914" target="\_blank">20491914</a>, PubMed:<a href="http://www.uniprot.org/citations/24790097" target="\_blank">24790097</a>). Ubiquitinates NFE2 by 'Lys-63' linkages and is implicated in the control of the development of hematopoietic lineages (PubMed:<a href="http://www.uniprot.org/citations/18718448" target="\_blank">18718448</a>). Mediates JUN ubiquitination and degradation (By similarity). Mediates JUNB ubiquitination and degradation (PubMed:<a href="http://www.uniprot.org/citations/16387660" target="\_blank">16387660</a>). Critical regulator of type 2 helper T (Th2) cell cytokine production by inducing JUNB ubiquitination and degradation (By similarity). Involved in the negative regulation of MAVS-dependent cellular antiviral responses (PubMed:<a href="http://www.uniprot.org/citations/19881509" target="\_blank">19881509</a>, PubMed:<a href="http://www.uniprot.org/citations/19881509" target="\_blank">19881509</a>). Ubiquitinates MAVS through 'Lys-48'-linked conjugation resulting in MAVS proteasomal degradation (PubMed:<a href="http://www.uniprot.org/citations/19881509" target="\_blank">19881509</a>). Following ligand stimulation, regulates sorting of Wnt receptor FZD4 to the degradative endocytic pathway probably by modulating PI42KA activity (PubMed:<a href="http://www.uniprot.org/citations/23146885" target="\_blank">23146885</a>). Ubiquitinates PI4K2A and negatively regulates its catalytic activity (PubMed:<a href="http://www.uniprot.org/citations/23146885" target="\_blank">23146885</a>). Ubiquitinates chemokine receptor CXCR4 and regulates sorting of CXCR4 to the degradative endocytic pathway following ligand stimulation by ubiquitinating endosomal sorting complex required for transport

ESCRT-0 components HGS and STAM (PubMed:<a href="http://www.uniprot.org/citations/14602072" target="\_blank">14602072</a>, PubMed:<a href="http://www.uniprot.org/citations/23146885" target="\_blank">23146885</a>, PubMed:<a href="http://www.uniprot.org/citations/34927784" target="\_blank">34927784</a>). Targets DTX1 for lysosomal degradation and controls NOTCH1 degradation, in the absence of ligand, through 'Lys-29'-linked polyubiquitination (PubMed:<a href="http://www.uniprot.org/citations/17028573" target="\_blank">17028573</a>, PubMed:<a href="http://www.uniprot.org/citations/18628966" target="\_blank">18628966</a>, PubMed:<a href="http://www.uniprot.org/citations/23886940" target="\_blank">23886940</a>). Ubiquitinates SNX9 (PubMed:<a href="http://www.uniprot.org/citations/20491914" target="\_blank">20491914</a>). Ubiquitinates MAP3K7 through 'Lys-48'-linked conjugation (By similarity). Together with UBR5, involved in the regulation of apoptosis and reactive oxygen species levels through the ubiquitination and proteasomal degradation of TXNIP: catalyzes 'Lys-48'-'Lys-63'-branched ubiquitination of TXNIP (PubMed:<a href="http://www.uniprot.org/citations/20068034" target="\_blank">20068034</a>, PubMed:<a href="http://www.uniprot.org/citations/29378950" target="\_blank">29378950</a>). ITCH synthesizes 'Lys-63'-linked chains, while UBR5 is branching multiple 'Lys-48'-linked chains of substrate initially modified (PubMed:<a href="http://www.uniprot.org/citations/29378950" target="\_blank">29378950</a>). Mediates the antiapoptotic activity of epidermal growth factor through the ubiquitination and proteasomal degradation of p15 BID (PubMed:<a href="http://www.uniprot.org/citations/20392206" target="\_blank">20392206</a>). Ubiquitinates BRAT1 and this ubiquitination is enhanced in the presence of NDFIP1 (PubMed:<a href="http://www.uniprot.org/citations/25631046" target="\_blank">25631046</a>). Inhibits the replication of influenza A virus (IAV) via ubiquitination of IAV matrix protein 1 (M1) through 'Lys-48'-linked conjugation resulting in M1 proteasomal degradation (PubMed:<a href="http://www.uniprot.org/citations/30328013" target="\_blank">30328013</a>). Ubiquitinates NEDD9/HEF1, resulting in proteasomal degradation of NEDD9/HEF1 (PubMed:<a href="http://www.uniprot.org/citations/15051726" target="\_blank">15051726</a>).

### Cellular Location

Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasm. Nucleus Early endosome membrane; Peripheral membrane protein; Cytoplasmic side. Endosome membrane; Peripheral membrane protein; Cytoplasmic side. Note=May be recruited to exosomes by NDFIP1 (PubMed:18819914). Localizes to plasma membrane upon CXCL12 stimulation where it co-localizes with CXCL4 (PubMed:14602072) Localization to early endosomes is increased upon CXCL12 stimulation where it co-localizes with DTX3L and CXCL4 (PubMed:24790097)

### Tissue Location

Widely expressed.

### ITCH Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### ITCH Antibody - Images



Western blot analysis of ITCH expression in K562 cell lysate.