

**Nuclear Matrix Protein p84 Antibody**  
**Rabbit mAb**  
**Catalog # AP91438****Specification**

---

**Nuclear Matrix Protein p84 Antibody - Product Information**

Application	WB, IHC, ICC
Primary Accession	<a href="#">Q96FV9</a>
Reactivity	Rat
Clonality	Monoclonal

**Other Names**

Death domain containing protein p84N5; HPR1; hTRES84; Nuclear matrix protein p84; p84N5; THO complex 1; Tho1; Thoc1;

Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	75666 Da

**Nuclear Matrix Protein p84 Antibody - Additional Information**

Dilution	WB~~1:1000 IHC~~1:100~500 ICC~~N/A
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Nuclear Matrix Protein p84
Description	Regulates transcriptional elongation of a subset of genes. Participates in an apoptotic pathway which is characterized by activation of caspase-6, increases in the expression of BAK1 and BCL2L1 and activation of NF-kappa-B.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

**Nuclear Matrix Protein p84 Antibody - Protein Information****Name** THOC1**Synonyms** HPR1**Function**

Component of the THO subcomplex of the TREX complex which is thought to couple mRNA transcription, processing and nuclear export, and which specifically associates with spliced mRNA and not with unspliced pre-mRNA (PubMed:<a href="http://www.uniprot.org/citations/15833825" target="\_blank">15833825</a>, PubMed:<a href="http://www.uniprot.org/citations/15998806"

target="\_blank">15998806</a>, PubMed:<a href="http://www.uniprot.org/citations/17190602" target="\_blank">17190602</a>). Required for efficient export of polyadenylated RNA (PubMed:<a href="http://www.uniprot.org/citations/23222130" target="\_blank">23222130</a>). The THOC1-THOC2-THOC3 core complex alone is sufficient to bind export factor NXF1-NXT1 and promote ATPase activity of DDX39B/UAP56 (PubMed:<a href="http://www.uniprot.org/citations/33191911" target="\_blank">33191911</a>). TREX is recruited to spliced mRNAs by a transcription-independent mechanism, binds to mRNA upstream of the exon-junction complex (EJC) and is recruited in a splicing- and cap- dependent manner to a region near the 5' end of the mRNA where it functions in mRNA export to the cytoplasm via the TAP/NXF1 pathway (PubMed:<a href="http://www.uniprot.org/citations/15833825" target="\_blank">15833825</a>, PubMed:<a href="http://www.uniprot.org/citations/15998806" target="\_blank">15998806</a>, PubMed:<a href="http://www.uniprot.org/citations/17190602" target="\_blank">17190602</a>). Regulates transcriptional elongation of a subset of genes (PubMed:<a href="http://www.uniprot.org/citations/22144908" target="\_blank">22144908</a>). Involved in genome stability by preventing co-transcriptional R-loop formation (By similarity). May play a role in hair cell formation, hence may be involved in hearing (By similarity).

### Cellular Location

[Isoform 1]: Nucleus speckle. Nucleus, nucleoplasm. Nucleus matrix. Cytoplasm. Note=Can shuttle between the nucleus and cytoplasm. Nuclear localization is required for induction of apoptotic cell death. Translocates to the cytoplasm during the early phase of apoptosis execution

### Tissue Location

Ubiquitous. Expressed in various cancer cell lines. Expressed at very low levels in normal breast epithelial cells and highly expressed in breast tumors. Expression is strongly associated with an aggressive phenotype of breast tumors and expression correlates with tumor size and the metastatic state of the tumor progression


## Nuclear Matrix Protein p84 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Nuclear Matrix Protein p84 Antibody - Images





Western blot analysis of Nuclear Matrix Protein p84 expression in HepG2 cell lysate.