

# FTO Antibody

Rabbit mAb Catalog # AP91635

### Specification

# FTO Antibody - Product Information

Application Primary Accession Clonality <b>Other Names</b> ALKBH9; Fto; GDFD; Protein fatso;	WB, IHC, ICC <u>09C0B1</u> Monoclonal
lsotype Host Calculated MW	Rabbit IgG Rabbit 58282 Da
FTO Antibody - Additional Information	
Dilution	WB~~1:1000 IHC~~1:100~500 ICC~~N/A
Purification Immunogen	Affinity-chromatography A synthesized peptide derived from human FTO
Description	Dioxygenase that repairs alkylated DNA and RNA by oxidative demethylation. Has highest activity towards single-stranded RNA containing 3-methyluracil, followed by single-stranded DNA containing 3-methylthymine.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

# **FTO Antibody - Protein Information**

Name FTO {ECO:0000303|PubMed:17496892, ECO:0000312|HGNC:HGNC:24678}

Function

RNA demethylase that mediates oxidative demethylation of different RNA species, such as mRNAs, tRNAs and snRNAs, and acts as a regulator of fat mass, adipogenesis and energy homeostasis (PubMed:<a href="http://www.uniprot.org/citations/22002720" target="\_blank">22002720</a>, PubMed:<a href="http://www.uniprot.org/citations/25452335" target="\_blank">25452335</a>, PubMed:<a href="http://www.uniprot.org/citations/26457839" target="\_blank">26457839</a>, PubMed:<a href="http://www.uniprot.org/citations/26457839" target="\_blank">26457839</a>, PubMed:<a href="http://www.uniprot.org/citations/26457839" target="\_blank">26457839</a>, PubMed:<a href="http://www.uniprot.org/citations/26457839" target="\_blank">26457839</a>, PubMed:<a href="http://www.uniprot.org/citations/26458103" target="\_blank">26458103</a>, PubMed:<a href="http://www.uniprot.org/citations/28002401" target="\_blank">26458103</a>, PubMed:<a href="http://www.uniprot.org/citations/28002401" target="\_blank">26458103</a>, PubMed:<a href="http://www.uniprot.org/citations/28002401" target="\_blank">26458103</a>, PubMed:<a href="http://www.uniprot.org/citations/30197295" target="\_blank">20197295</a>).



Specifically demethylates N(6)- methyladenosine (m6A) RNA, the most prevalent internal modification of messenger RNA (mRNA) in higher eukaryotes (PubMed:<a href="http://www.uniprot.org/citations/22002720" target=" blank">22002720</a>, PubMed:<a href="http://www.uniprot.org/citations/25452335" target="\_blank">25452335</a>, PubMed:<a href="http://www.uniprot.org/citations/26457839" target=" blank">26457839</a>, PubMed:<a href="http://www.uniprot.org/citations/26458103" target=" blank">26458103</a>, PubMed:<a href="http://www.uniprot.org/citations/30197295" target=" blank">30197295</a>). M6A demethylation by FTO affects mRNA expression and stability (PubMed:<a href="http://www.uniprot.org/citations/30197295" target=" blank">30197295</a>). Also able to demethylate m6A in U6 small nuclear RNA (snRNA) (PubMed:<a href="http://www.uniprot.org/citations/30197295" target=" blank">30197295</a>). Mediates demethylation of N(6),2'-O- dimethyladenosine cap (m6A(m)), by demethylating the N(6)methyladenosine at the second transcribed position of mRNAs and U6 snRNA (PubMed:<a href="http://www.uniprot.org/citations/28002401" target=" blank">28002401</a>, PubMed:<a href="http://www.uniprot.org/citations/30197295" target=" blank">30197295</a>). Demethylation of m6A(m) in the 5'-cap by FTO affects mRNA stability by promoting susceptibility to decapping (PubMed:<a href="http://www.uniprot.org/citations/28002401" target=" blank">28002401</a>). Also acts as a tRNA demethylase by removing N(1)-methyladenine from various tRNAs (PubMed:<a href="http://www.uniprot.org/citations/30197295" target=" blank">30197295</a>). Has no activity towards 1-methylguanine (PubMed: <a href="http://www.uniprot.org/citations/20376003" target=" blank">20376003</a>). Has no detectable activity towards double-stranded DNA (PubMed:<a href="http://www.uniprot.org/citations/20376003" target=" blank">20376003</a>). Also able to repair alkylated DNA and RNA by oxidative demethylation: demethylates single-stranded RNA containing 3-methyluracil, single- stranded DNA containing 3-methylthymine and has low demethylase activity towards single-stranded DNA containing 1-methyladenine or 3methylcytosine (PubMed:<a href="http://www.uniprot.org/citations/18775698" target=" blank">18775698</a>, PubMed:<a href="http://www.uniprot.org/citations/20376003" target=" blank">20376003</a>). Ability to repair alkylated DNA and RNA is however unsure in vivo (PubMed:<a href="http://www.uniprot.org/citations/18775698" target=" blank">18775698</a>, PubMed:<a href="http://www.uniprot.org/citations/20376003" target=" blank">20376003</a>). Involved in the regulation of fat mass, adipogenesis and body weight, thereby contributing to the regulation of body size and body fat accumulation (PubMed:<a href="http://www.uniprot.org/citations/18775698" target=" blank">18775698</a>, PubMed:<a href="http://www.uniprot.org/citations/20376003" target=" blank">20376003</a>). Involved in the regulation of thermogenesis and the control of adipocyte differentiation into brown or white fat cells (PubMed:<a href="http://www.uniprot.org/citations/26287746" target=" blank">26287746</a>). Regulates activity of the dopaminergic midbrain circuitry via its ability to demethylate m6A in mRNAs (By similarity). Plays an oncogenic role in a number of acute myeloid leukemias by enhancing leukemic oncogene-mediated cell transformation: acts by mediating m6A demethylation of target transcripts such as MYC, CEBPA, ASB2 and RARA, leading to promote their expression (PubMed:<a href="http://www.uniprot.org/citations/28017614" target=" blank">28017614</a>, PubMed:<a href="http://www.uniprot.org/citations/29249359" target=" blank">29249359</a>).

#### **Cellular Location**

Nucleus. Nucleus speckle. Cytoplasm Note=Localizes mainly in the nucleus, where it is able to demethylate N(6)-methyladenosine (m6A) and N(6),2'-O-dimethyladenosine cap (m6A(m)) in U6 small nuclear RNA (snRNA), N(1)-methyladenine from tRNAs and internal m6A in mRNAs (PubMed:30197295). In the cytoplasm, mediates demethylation of m6A and m6A(m) in mRNAs and N(1)-methyladenine from tRNAs (PubMed:30197295).

#### **Tissue Location**

Ubiquitously expressed, with relatively high expression in adrenal glands and brain; especially in hypothalamus and pituitary (PubMed:17434869, PubMed:17496892). Highly expressed in highly expressed in acute myeloid leukemias (AML) with t(11;11)(q23;23) with KMT2A/MLL1 rearrangements, t(15;17)(q21;q21)/PML-RARA, FLT3-ITD, and/or NPM1 mutations



(PubMed:28017614).

### FTO Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

### FTO Antibody - Images



Western blot analysis of FTO expression in HEK293 cell lysate.