

### **TXNIP Antibody**

Rabbit mAb Catalog # AP91646

# **Specification**

### **TXNIP Antibody - Product Information**

Application WB, IHC, FC, ICC

Primary Accession
Reactivity
Q9H3M7
Rat

Clonality Monoclonal

**Other Names** 

HHCPA78; THIF; TXNIP; VDUP1;

Isotype Rabbit IgG
Host Rabbit
Calculated MW 43661 Da

## **TXNIP Antibody - Additional Information**

Dilution WB~~1:1000

IHC~~1:100~500 FC~~1:10~50 ICC~~N/A

Purification Affinity-chromatography

Immunogen A synthesized peptide derived from human

**TXNIP** 

Description May act as an oxidative stress mediator by

inhibiting thioredoxin activity or by limiting its bioavailability. Interacts with COPS5 and restores COPS5-induced suppression of CDKN1B stability, blocking the COPS5-mediated translocation of

CDKN1B from the nucleus to the

cytoplasm.

Storage Condition and Buffer Rabbit IgG in phosphate buffered saline ,

pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid

freeze / thaw cycle.

# **TXNIP Antibody - Protein Information**

**Name TXNIP** 

Synonyms VDUP1

#### **Function**

May act as an oxidative stress mediator by inhibiting thioredoxin activity or by limiting its bioavailability (PubMed:<a href="http://www.uniprot.org/citations/17603038"">http://www.uniprot.org/citations/17603038</a>"



target="\_blank">17603038</a>). Interacts with COPS5 and restores COPS5-induced suppression of CDKN1B stability, blocking the COPS5-mediated translocation of CDKN1B from the nucleus to the cytoplasm (By similarity). Functions as a transcriptional repressor, possibly by acting as a bridge molecule between transcription factors and corepressor complexes, and over-expression will induce G0/G1 cell cycle arrest (PubMed:<a href="http://www.uniprot.org/citations/12821938" target="\_blank">12821938</a>). Required for the maturation of natural killer cells (By similarity). Acts as a suppressor of tumor cell growth (PubMed:<a

href="http://www.uniprot.org/citations/18541147" target="\_blank">18541147</a>). Inhibits the proteasomal degradation of DDIT4, and thereby contributes to the inhibition of the mammalian target of rapamycin complex 1 (mTORC1) (PubMed:<a

href="http://www.uniprot.org/citations/21460850" target=" blank">21460850</a>).


**Cellular Location** Cytoplasm. Nucleus

## **TXNIP Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

## **TXNIP Antibody - Images**



Western blot analysis of TXNIP expression in BxPC-3 cell lysate.