

**HAUSP / USP7 Antibody  
Rabbit mAb  
Catalog # AP91649**

## Specification

## **HAUSP / USP7 Antibody - Product Information**

Application	WB, IHC, FC, ICC
Primary Accession	<a href="#">Q93009</a>
Reactivity	Rat
Clonality	Monoclonal
<b>Other Names</b>	
TEF1; HAUSP; USP7;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	128302 Da

## HAUSP / USP7 Antibody - Additional Information

Dilution	WB~~1:1000 IHC~~1:100~500 FC~~1:10~50 ICC~~N/A
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human HAUSP / USP7
Description	Hydrolase that deubiquitinates target proteins such as FOXO4, p53/TP53, MDM2, ERCC6, DNMT1, UHRF1, PTEN and DAXX (PubMed:11923872, PubMed:15053880, PubMed:16964248, PubMed:18716620, PubMed:25283148).
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

## HAUSP / USP7 Antibody - Protein Information

**Name** USP7 {ECO:0000303|PubMed:12093161, ECO:0000312|HGNC:HGNC:12630}

## Function

Hydrolase that deubiquitinates target proteins such as ARMC5, FOXO4, DEPTOR, KAT5, p53/TP53, MDM2, ERCC6, DNMT1, UHRF1, PTEN, KMT2E/MLL5 and DAXX (PubMed:[11923872](http://www.uniprot.org/citations/11923872), PubMed:[15053880](http://www.uniprot.org/citations/15053880), PubMed:[16964248](http://www.uniprot.org/citations/16964248), PubMed:[18716620](http://www.uniprot.org/citations/18716620))

href="http://www.uniprot.org/citations/25283148" target="\_blank">>25283148</a>, PubMed:<a href="http://www.uniprot.org/citations/25865756" target="\_blank">>25865756</a>, PubMed:<a href="http://www.uniprot.org/citations/26678539" target="\_blank">>26678539</a>, PubMed:<a href="http://www.uniprot.org/citations/28655758" target="\_blank">>28655758</a>, PubMed:<a href="http://www.uniprot.org/citations/33544460" target="\_blank">>33544460</a>, PubMed:<a href="http://www.uniprot.org/citations/35216969" target="\_blank">>35216969</a>). Together with DAXX, prevents MDM2 self-ubiquitination and enhances the E3 ligase activity of MDM2 towards p53/TP53, thereby promoting p53/TP53 ubiquitination and proteasomal degradation (PubMed:<a href="http://www.uniprot.org/citations/15053880" target="\_blank">>15053880</a>, PubMed:<a href="http://www.uniprot.org/citations/16845383" target="\_blank">>16845383</a>, PubMed:<a href="http://www.uniprot.org/citations/18566590" target="\_blank">>18566590</a>, PubMed:<a href="http://www.uniprot.org/citations/20153724" target="\_blank">>20153724</a>). Deubiquitinates p53/TP53, preventing degradation of p53/TP53, and enhances p53/TP53-dependent transcription regulation, cell growth repression and apoptosis (PubMed:<a href="http://www.uniprot.org/citations/25283148" target="\_blank">>25283148</a>). Deubiquitinates p53/TP53 and MDM2 and strongly stabilizes p53/TP53 even in the presence of excess MDM2, and also induces p53/TP53-dependent cell growth repression and apoptosis (PubMed:<a href="http://www.uniprot.org/citations/11923872" target="\_blank">>11923872</a>, PubMed:<a href="http://www.uniprot.org/citations/26786098" target="\_blank">>26786098</a>). Deubiquitination of FOXO4 in presence of hydrogen peroxide is not dependent on p53/TP53 and inhibits FOXO4-induced transcriptional activity (PubMed:<a href="http://www.uniprot.org/citations/16964248" target="\_blank">>16964248</a>). In association with DAXX, is involved in the deubiquitination and translocation of PTEN from the nucleus to the cytoplasm, both processes that are counteracted by PML (PubMed:<a href="http://www.uniprot.org/citations/18716620" target="\_blank">>18716620</a>). Deubiquitinates KMT2E/MLL5 preventing KMT2E/MLL5 proteasomal-mediated degradation (PubMed:<a href="http://www.uniprot.org/citations/26678539" target="\_blank">>26678539</a>). Involved in cell proliferation during early embryonic development. Involved in transcription-coupled nucleotide excision repair (TC-NER) in response to UV damage: recruited to DNA damage sites following interaction with KIAA1530/UVSSA and promotes deubiquitination of ERCC6, preventing UV-induced degradation of ERCC6 (PubMed:<a href="http://www.uniprot.org/citations/22466611" target="\_blank">>22466611</a>, PubMed:<a href="http://www.uniprot.org/citations/22466612" target="\_blank">>22466612</a>). Involved in maintenance of DNA methylation via its interaction with UHRF1 and DNMT1: acts by mediating deubiquitination of UHRF1 and DNMT1, preventing their degradation and promoting DNA methylation by DNMT1 (PubMed:<a href="http://www.uniprot.org/citations/21745816" target="\_blank">>21745816</a>, PubMed:<a href="http://www.uniprot.org/citations/22411829" target="\_blank">>22411829</a>). Deubiquitinates alkylation repair enzyme ALKBH3. OTUD4 recruits USP7 and USP9X to stabilize ALKBH3, thereby promoting the repair of alkylated DNA lesions (PubMed:<a href="http://www.uniprot.org/citations/25944111" target="\_blank">>25944111</a>). Acts as a chromatin regulator via its association with the Polycomb group (PcG) multiprotein PRC1-like complex; may act by deubiquitinating components of the PRC1-like complex (PubMed:<a href="http://www.uniprot.org/citations/20601937" target="\_blank">>20601937</a>). Able to mediate deubiquitination of histone H2B; it is however unsure whether this activity takes place in vivo (PubMed:<a href="http://www.uniprot.org/citations/20601937" target="\_blank">>20601937</a>). Exhibits a preference towards 'Lys-48'-linked ubiquitin chains (PubMed:<a href="http://www.uniprot.org/citations/22689415" target="\_blank">>22689415</a>). Increases regulatory T-cells (Treg) suppressive capacity by deubiquitinating and stabilizing the transcription factor FOXP3 which is crucial for Treg cell function (PubMed:<a href="http://www.uniprot.org/citations/23973222" target="\_blank">>23973222</a>). Plays a role in the maintenance of the circadian clock periodicity via deubiquitination and stabilization of the CRY1 and CRY2 proteins (PubMed:<a href="http://www.uniprot.org/citations/27123980" target="\_blank">>27123980</a>). Deubiquitinates REST, thereby stabilizing REST and promoting the maintenance of neural progenitor cells (PubMed:<a href="http://www.uniprot.org/citations/21258371" target="\_blank">>21258371</a>). Deubiquitinates SIRT7, inhibiting SIRT7 histone deacetylase activity and regulating

gluconeogenesis (PubMed:<a href="http://www.uniprot.org/citations/28655758" target="\_blank">28655758</a>). Involved in the regulation of WASH-dependent actin polymerization at the surface of endosomes and the regulation of endosomal protein recycling (PubMed:<a href="http://www.uniprot.org/citations/26365382" target="\_blank">26365382</a>). It maintains optimal WASH complex activity and precise F-actin levels via deubiquitination of TRIM27 and WASHC1 (PubMed:<a href="http://www.uniprot.org/citations/26365382" target="\_blank">26365382</a> target="\_blank">26365382</a>). Mediates the deubiquitination of phosphorylated DEPTOR, promoting its stability and leading to decreased mTORC1 signaling (PubMed:<a href="http://www.uniprot.org/citations/35216969" target="\_blank">35216969</a>).

#### Cellular Location

Nucleus. Cytoplasm Nucleus. PML body. Chromosome. Note=Present in a minority of ND10 nuclear bodies. Association with ICP0/VMW110 at early times of infection leads to an increased proportion of USP7-containing ND10 Colocalizes with ATXN1 in the nucleus. Colocalized with DAXX in speckled structures. Colocalized with PML and PTEN in promyelocytic leukemia protein (PML) nuclear bodies

#### Tissue Location


Expressed in neural progenitor cells (at protein level) (PubMed:21258371). Widely expressed. Overexpressed in prostate cancer.

### HAUSP / USP7 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### HAUSP / USP7 Antibody - Images



Western blot analysis of HAUSP / USP7 expression in Ramos cell lysate.