

CXCL11 Antibody

Rabbit mAb Catalog # AP91704

Specification

CXCL11 Antibody - Product Information

Application WB, ICC
Primary Accession O14625
Clonality Monoclonal

Other Names

betaR; CXC11; CXCL11; H174; I TAC; IP9; ITAC; SCYB11; SCYB9B;

Isotype Rabbit IgG
Host Rabbit
Calculated MW 10365 Da

CXCL11 Antibody - Additional Information

Dilution WB~~1:1000

ICC~~N/A

Purification Affinity-chromatography

Immunogen A synthesized peptide derived from human

CXCL11

Description Chemotactic for interleukin-activated

T-cells but not unstimulated T-cells,

neutrophils or monocytes. Induces calcium

release in activated T-cells. Binds to

CXCR3. May play an important role in CNS diseases which involve T-cell recruitment. May play a role in skin immune responses. Rabbit IgG in phosphate buffered saline,

pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid

freeze / thaw cycle.

CXCL11 Antibody - Protein Information

Name CXCL11

Synonyms ITAC, SCYB11, SCYB9B

Storage Condition and Buffer

Function

Chemotactic for interleukin-activated T-cells but not unstimulated T-cells, neutrophils or monocytes. Induces calcium release in activated T-cells. Binds to CXCR3. May play an important role in CNS diseases which involve T-cell recruitment. May play a role in skin immune responses.

Cellular Location

Secreted.





Tissue Location


High levels in peripheral blood leukocytes, pancreas and liver astrocytes. Moderate levels in thymus, spleen and lung. Low levels in placenta, prostate and small intestine. Also found in epidermal basal layer keratinocytes in skin disorders

CXCL11 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

CXCL11 Antibody - Images



Western blot analysis of CXCL11 expression in THP1 cell lysate.