

**CYP26A1 Antibody**  
**Rabbit mAb**  
**Catalog # AP91710****Specification****CYP26A1 Antibody - Product Information**

Application	WB, IHC
Primary Accession	<a href="#">O43174</a>
Reactivity	Rat
Clonality	Monoclonal
<b>Other Names</b>	
CP26; CYP26; cyp26a1; hP450RAI; P450RAI; P450RAI1;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	56199 Da

**CYP26A1 Antibody - Additional Information**

Dilution	WB~~1:1000 IHC~~1:100~500
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human CYP26A1
Description	Plays a key role in retinoic acid metabolism. Acts on retinoids, including all-trans-retinoic acid (RA) and its stereoisomer 9-cis-RA. Capable of both 4-hydroxylation and 18-hydroxylation. Responsible for generation of several hydroxylated forms of RA, including 4-OH-RA, 4-oxo-RA and 18-OH-RA.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

**CYP26A1 Antibody - Protein Information**

**Name** CYP26A1 {ECO:0000303|PubMed:26937021, ECO:0000312|HGNC:HGNC:2603}

**Function**

A cytochrome P450 monooxygenase involved in the metabolism of retinoates (RAs), the active metabolites of vitamin A, and critical signaling molecules in animals (PubMed:<a href="http://www.uniprot.org/citations/22020119" target="\_blank">22020119</a>, PubMed:<a href="http://www.uniprot.org/citations/9228017" target="\_blank">9228017</a>, PubMed:<a href="http://www.uniprot.org/citations/9716180" target="\_blank">9716180</a>). RAs exist as at least four different isomers: all- trans-RA (atRA), 9-cis-RA, 13-cis-RA, and 9,13-dicis-RA, where atRA

is considered to be the biologically active isomer, although 9-cis-RA and 13-cis-RA also have activity (Probable). Catalyzes the hydroxylation of atRA primarily at C-4 and C-18, thereby contributing to the regulation of atRA homeostasis and signaling (PubMed:<a href="http://www.uniprot.org/citations/22020119" target="\_blank">22020119</a>, PubMed:<a href="http://www.uniprot.org/citations/9228017" target="\_blank">9228017</a>, PubMed:<a href="http://www.uniprot.org/citations/9716180" target="\_blank">9716180</a>). Hydroxylation of atRA limits its biological activity and initiates a degradative process leading to its eventual elimination (Probable). Involved in the conversion of atRA to all-trans-4-oxo-RA. Able to metabolize other RAs such as 9-cis, 13-cis and 9,13-di-cis RA (By similarity) (PubMed:<a href="http://www.uniprot.org/citations/9228017" target="\_blank">9228017</a>). Can oxidize all-trans-13,14- dihydroretinoate (DRA) to metabolites which could include all-trans-4- oxo-DRA, all-trans-4-hydroxy-DRA, all-trans-5,8-epoxy-DRA, and all- trans-18-hydroxy-DRA (By similarity). May play a role in the oxidative metabolism of xenobiotics such as tazarotenic acid (PubMed:<a href="http://www.uniprot.org/citations/26937021" target="\_blank">26937021</a>).

#### **Cellular Location**

Endoplasmic reticulum membrane; Peripheral membrane protein. Microsome membrane; Peripheral membrane protein

#### **Tissue Location**

Expressed in most fetal and adult tissues with highest levels in adult liver, heart, pituitary gland, adrenal gland, placenta and regions of the brain (PubMed:9826557). Expressed at high levels in lung, pancreas, skin and uterus (at protein level) (PubMed:22020119). Lower expression level is detected in spleen, kidney, intestine and adipose tissue (at protein level) (PubMed:22020119).


### **CYP26A1 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **CYP26A1 Antibody - Images**





Western blot analysis of CYP26A1 expression in HepG2 cell lysate.