

**GRASP65 Antibody**  
**Rabbit mAb**  
**Catalog # AP91729****Specification**

---

**GRASP65 Antibody - Product Information**

|                           |                        |
|---------------------------|------------------------|
| Application               | WB, IHC, FC, ICC, IP   |
| Primary Accession         | <a href="#">O9BQQ3</a> |
| Clonality                 | Monoclonal             |
| <b>Other Names</b>        |                        |
| GOLPH5; GORASP1; GRASP65; |                        |
| Isotype                   | Rabbit IgG             |
| Host                      | Rabbit                 |
| Calculated MW             | 46482 Da               |

**GRASP65 Antibody - Additional Information**

|                              |  |
|------------------------------|--|
| Dilution                     | WB~~1:1000<br>IHC~~1:100~500<br>FC~~1:10~50<br>ICC~~N/A<br>IP~~N/A   |
| Purification                 | Affinity-chromatography  |
| Immunogen                    | A synthesized peptide derived from human GRASP65   |
| Description                  | Stacking factor involved in the postmitotic assembly of Golgi stacks from mitotic Golgi fragments. Key structural protein required for the maintenance of the Golgi apparatus integrity: its caspase-mediated cleavage is required for fragmentation of the Golgi during apoptosis |
| Storage Condition and Buffer | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.  |

**GRASP65 Antibody - Protein Information****Name** GORASP1**Synonyms** GOLPH5, GRASP65**Function**

Key structural protein of the Golgi apparatus (PubMed:<a href="http://www.uniprot.org/citations/33301566" target="\_blank">33301566</a>). The membrane cisternae of the Golgi apparatus adhere to each other to form stacks, which are aligned

side by side to form the Golgi ribbon (PubMed:<a href="http://www.uniprot.org/citations/33301566" target="\_blank">33301566</a>). Acting in concert with GORASP2/GRASP55, is required for the formation and maintenance of the Golgi ribbon, and may be dispensable for the formation of stacks (PubMed:<a href="http://www.uniprot.org/citations/33301566" target="\_blank">33301566</a>). However, other studies suggest that GORASP1 plays an important role in assembly and membrane stacking of the cisternae, and in the reassembly of Golgi stacks after breakdown during mitosis (By similarity). Caspase-mediated cleavage of GORASP1 is required for fragmentation of the Golgi during apoptosis (By similarity). Also mediates, via its interaction with GOLGA2/GM130, the docking of transport vesicles with the Golgi membranes (PubMed:<a href="http://www.uniprot.org/citations/16489344" target="\_blank">16489344</a>). Mediates ER stress-induced unconventional (ER/Golgi-independent) trafficking of core-glycosylated CFTR to cell membrane (PubMed:<a href="http://www.uniprot.org/citations/21884936" target="\_blank">21884936</a>).

### Cellular Location


Golgi apparatus, cis-Golgi network membrane; Peripheral membrane protein; Cytoplasmic side. Endoplasmic reticulum- Golgi intermediate compartment membrane

### GRASP65 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### GRASP65 Antibody - Images



Western blot analysis of GRASP65 expression in MCF7 cell lysate.