

APPL Antibody
Rabbit mAb
Catalog # AP91758**Specification**

APPL Antibody - Product Information

Application	WB, IHC, ICC
Primary Accession	O9UKG1
Reactivity	Rat
Clonality	Monoclonal
Other Names	
DIP13 alpha; DP13A; KIAA1428; APPL1; Dip13-alpha; Adapter protein containing PH domain; DCC-interacting protein 13-alpha;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	79663 Da

APPL Antibody - Additional Information

Dilution	WB~~1:1000 IHC~~1:100~500 ICC~~N/A
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human APPL
Description	Adapter protein that interacts with proteins involved in different cellular signaling pathways. Required for the regulation of cell proliferation in response to extracellular signals from an early endosomal compartment. Links Rab5 to nuclear signal transduction. Involved in the regulation of the insulin receptor signaling pathway.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

APPL Antibody - Protein Information**Name** APPL1 ([HGNC:24035](#))**Function**

Multifunctional adapter protein that binds to various membrane receptors, nuclear factors and signaling proteins to regulate many processes, such as cell proliferation, immune response, endosomal trafficking and cell metabolism (PubMed:<a

[10490823](http://www.uniprot.org/citations/10490823), PubMed: [15016378](http://www.uniprot.org/citations/15016378), PubMed: [19661063](http://www.uniprot.org/citations/19661063), PubMed: [26073777](http://www.uniprot.org/citations/26073777), PubMed: [26583432](http://www.uniprot.org/citations/26583432)). Regulates signaling pathway leading to cell proliferation through interaction with RAB5A and subunits of the NuRD/MeCP1 complex (PubMed: [15016378](http://www.uniprot.org/citations/15016378)). Functions as a positive regulator of innate immune response via activation of AKT1 signaling pathway by forming a complex with APPL1 and PIK3R1 (By similarity). Inhibits Fc-gamma receptor-mediated phagocytosis through PI3K/Akt signaling in macrophages (By similarity). Regulates TLR4 signaling in activated macrophages (By similarity). Involved in trafficking of the TGFBR1 from the endosomes to the nucleus via microtubules in a TRAF6-dependent manner (PubMed: [26583432](http://www.uniprot.org/citations/26583432)). Plays a role in cell metabolism by regulating adiponectin and insulin signaling pathways (PubMed: [19661063](http://www.uniprot.org/citations/19661063), PubMed: [24879834](http://www.uniprot.org/citations/24879834), PubMed: [26073777](http://www.uniprot.org/citations/26073777)). Required for fibroblast migration through HGF cell signaling (By similarity). Positive regulator of beta-catenin/TCF-dependent transcription through direct interaction with RUVBL2/reptin resulting in the relief of RUVBL2-mediated repression of beta-catenin/TCF target genes by modulating the interactions within the beta-catenin-reptin-HDAC complex (PubMed: [19433865](http://www.uniprot.org/citations/19433865)).

Cellular Location

Early endosome membrane; Peripheral membrane protein. Nucleus. Cytoplasm. Endosome. Cell projection, ruffle {ECO:0000250|UniProtKB:Q8K3H0}. Cytoplasmic vesicle, phagosome {ECO:0000250|UniProtKB:Q8K3H0}. Note=Early endosomal membrane-bound and nuclear. Translocated into the nucleus upon release from endosomal membranes following internalization of EGF

Tissue Location


High levels in heart, ovary, pancreas and skeletal muscle.


APPL Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

APPL Antibody - Images





Western blot analysis of APPL expression in HeLa cell lysate.