

CHD1L Antibody
Rabbit mAb
Catalog # AP91778**Specification**

CHD1L Antibody - Product Information

Application	WB, IHC, FC, ICC, IP
Primary Accession	Q86WJ1
Clonality	Monoclonal
Other Names	
ALC1; chd1l; CHDL;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	101000 Da

CHD1L Antibody - Additional Information

Dilution	WB~~1:1000 IHC~~1:100~500 FC~~1:10~50 ICC~~N/A IP~~N/A
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human CHD1L
Description	DNA helicase which plays a role in chromatin-remodeling following DNA damage. Targeted to sites of DNA damage through interaction with poly(ADP-ribose) and functions to regulate chromatin during DNA repair. Able to catalyze nucleosome sliding in an ATP-dependent manner. Helicase activity is strongly stimulated upon poly(ADP-ribose)-binding.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

CHD1L Antibody - Protein Information**Name** CHD1L {ECO:0000303|PubMed:34210977, ECO:0000312|HGNC:HGNC:1916}**Function**ATP-dependent chromatin remodeler that mediates chromatin- remodeling following DNA damage (PubMed:19661379,
PubMed:29220652,

PubMed:29220653, PubMed:33357431, PubMed:34210977, PubMed:34486521, PubMed:34874266). Recruited to DNA damage sites through interaction with poly-ADP-ribose: specifically recognizes and binds histones that are poly-ADP-ribosylated on serine residues in response to DNA damage (PubMed:19661379, PubMed:29220652, PubMed:29220653, PubMed:34486521, PubMed:34874266). Poly-ADP-ribose-binding activates the ATP-dependent chromatin remodeler activity, thereby regulating chromatin during DNA repair (PubMed:19661379, PubMed:29220652, PubMed:29220653, PubMed:34486521, PubMed:34874266). Catalyzes nucleosome sliding away from DNA breaks in an ATP-dependent manner (PubMed:19661379, PubMed:29220652, PubMed:29220653). Chromatin remodeling activity promotes PARP2 removal from chromatin (PubMed:33275888).

Cellular Location

Nucleus. Chromosome Note=Localizes at sites of DNA damage; recruited by histones H2B and H3 poly-ADP-ribosylated on 'Ser-6' and 'Ser-10', respectively (H2BS6ADPr and H3S10ADPr) by PARP1 or PARP2.

Tissue Location


Frequently overexpressed in hepatomacellular carcinomas.


CHD1L Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CHD1L Antibody - Images





Western blot analysis of CHD1L expression in A549 cell lysate.