

KAT13A / SRC1 Antibody
Rabbit mAb
Catalog # AP91828**Specification**

KAT13A / SRC1 Antibody - Product Information

Application	WB, IHC, IP
Primary Accession	Q15788
Clonality	Monoclonal
Other Names	
bHLHe74; mNRC 1; NCoA 1; RIP160; SRC 1;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	156757 Da

KAT13A / SRC1 Antibody - Additional Information

Dilution	WB~~1:1000 IHC~~1:100~500 IP~~N/A
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human KAT13A / SRC1
Description	Nuclear receptor coactivator that directly binds nuclear receptors and stimulates the transcriptional activities in a hormone-dependent fashion. Involved in the coactivation of different nuclear receptors, such as for steroids (PGR, GR and ER), retinoids (RXRs), thyroid hormone (TRs) and prostanoids (PPARs).
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

KAT13A / SRC1 Antibody - Protein Information**Name** NCOA1**Synonyms** BHLHE74, SRC1**Function**

Nuclear receptor coactivator that directly binds nuclear receptors and stimulates the transcriptional activities in a hormone- dependent fashion. Involved in the coactivation of different nuclear receptors, such as for steroids (PGR, GR and ER), retinoids (RXRs), thyroid hormone (TRs) and prostanoids (PPARs). Also involved in coactivation mediated by STAT3, STAT5A, STAT5B and

STAT6 transcription factors. Displays histone acetyltransferase activity toward H3 and H4; the relevance of such activity remains however unclear. Plays a central role in creating multisubunit coactivator complexes that act via remodeling of chromatin, and possibly acts by participating in both chromatin remodeling and recruitment of general transcription factors. Required with NCOA2 to control energy balance between white and brown adipose tissues. Required for mediating steroid hormone response. Isoform 2 has a higher thyroid hormone-dependent transactivation activity than isoform 1 and isoform 3.

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00981}.


Tissue Location

Widely expressed.

KAT13A / SRC1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

KAT13A / SRC1 Antibody - Images

Western blot analysis of KAT13A / SRC1 expression in HEK293 cell lysate.