

**MC4 Receptor Antibody**  
**Rabbit mAb**  
**Catalog # AP91945****Specification**

---

**MC4 Receptor Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">P32245</a>
Reactivity	Rat
Clonality	Monoclonal
<b>Other Names</b>	
MC4R; Melanocortin 4 receptor;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	36943 Da

**MC4 Receptor Antibody - Additional Information**

Dilution	WB~~1:1000
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human MC4 Receptor
Description	Receptor specific to the heptapeptide core common to adrenocorticotrophic hormone and alpha-, beta-, and gamma-MSH. This receptor is mediated by G proteins that stimulate adenylate cyclase.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

**MC4 Receptor Antibody - Protein Information****Name** MC4R**Function**

Hormone receptor that acts as a key component of the leptin- melanocortin pathway at the intersection of homeostatic maintenance of energetic state (PubMed:<a href="http://www.uniprot.org/citations/32327598" target="\_blank">32327598</a>, PubMed:<a href="http://www.uniprot.org/citations/33858992" target="\_blank">33858992</a>). Plays a role in regulating food intake: activation by a stimulating hormone such as anorexigenic alpha-melanocyte stimulating hormone (alpha-MSH) inhibits appetite, whereas binding to a natural antagonist like Agouti-related protein/AGRP promotes appetite. G-protein-coupled receptor that activates conventional G $\alpha$ s signaling leading to induction of anorexigenic signaling in the hypothalamus to result in negative energy balance (PubMed:<a href="http://www.uniprot.org/citations/33858992" target="\_blank">33858992</a>). Regulates

the firing activity of neurons from the hypothalamus by alpha-MSH and AGRP independently of Galphas signaling by ligand-induced coupling of closure of inwardly rectifying potassium channel KCNJ13 (By similarity). In intestinal epithelial cells, plays a role in the inhibition of hepatic glucose production via nesfatin-1/NUCB2 leading to increased cyclic adenosine monophosphate (cAMP) levels and glucagon-like peptide 1 (GLP-1) secretion in the intestinal epithelium (PubMed:<a href="http://www.uniprot.org/citations/39562740" target="\_blank">39562740</a>).

#### Cellular Location

Cell membrane; Multi-pass membrane protein

#### Tissue Location


Brain, placental, and gut tissues.

### MC4 Receptor Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### MC4 Receptor Antibody - Images



Western blot analysis of MC4 Receptor expression in (1) MCF7 cell lysate; (2) RAW264.7 cell lysate.