

EAAT3 Antibody
Rabbit mAb
Catalog # AP91955**Specification**

EAAT3 Antibody - Product Information

Application	WB, IHC, ICC
Primary Accession	P43005
Reactivity	Rat
Clonality	Monoclonal
Other Names	
SLC1A1, EAAC1, EAAT3, Eaac-1;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	57100 Da

EAAT3 Antibody - Additional Information

Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human EAAT3
Description	Transports L-glutamate and also L- and D-aspartate. Essential for terminating the postsynaptic action of glutamate by rapidly removing released glutamate from the synaptic cleft. Acts as a symport by cotransporting sodium. Negatively regulated by ARL6IP5.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

EAAT3 Antibody - Protein Information**Name** SLC1A1 ([HGNC:10939](#))**Function**

Sodium-dependent, high-affinity amino acid transporter that mediates the uptake of L-glutamate and also L-aspartate and D-aspartate (PubMed:21123949, PubMed:26690923, PubMed:33658209, PubMed:7521911, PubMed:7914198, PubMed:8857541). Can also transport L-cysteine (PubMed:<a href="http://www.uniprot.org/citations/21123949"

target="_blank">21123949). Functions as a symporter that transports one amino acid molecule together with two or three Na(+) ions and one proton, in parallel with the counter-transport of one K(+) ion (PubMed:26690923, PubMed:33658209, PubMed:7521911, PubMed:8857541). Mediates Cl(-) flux that is not coupled to amino acid transport; this avoids the accumulation of negative charges due to aspartate and Na(+) symport (PubMed:26690923, PubMed:8857541). Plays an important role in L- glutamate and L-aspartate reabsorption in renal tubuli (PubMed:21123949). Plays a redundant role in the rapid removal of released glutamate from the synaptic cleft, which is essential for terminating the postsynaptic action of glutamate (By similarity). Contributes to glutathione biosynthesis and protection against oxidative stress via its role in L-glutamate and L-cysteine transport (By similarity). Negatively regulated by ARL6IP5 (By similarity).

Cellular Location

Cell membrane; Multi-pass membrane protein {ECO:0000250|UniProtKB:P43003}. Apical cell membrane; Multi-pass membrane protein {ECO:0000250|UniProtKB:P43003}. Synapse, synaptosome {ECO:0000250|UniProtKB:P51906}. Early endosome membrane {ECO:0000250|UniProtKB:P51906}. Late endosome membrane {ECO:0000250|UniProtKB:P51906}. Recycling endosome membrane {ECO:0000250|UniProtKB:P51906}

Tissue Location


Expressed in all tissues tested including liver, muscle, testis, ovary, retinoblastoma cell line, neurons and brain (in which there was dense expression in substantia nigra, red nucleus, hippocampus and in cerebral cortical layers)


EAAT3 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

EAAT3 Antibody - Images





Western blot analysis of EAAT3 expression in Human fetal brain lysate.