

### FBXO3 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP9195b

## **Specification**

#### FBXO3 Antibody (C-term) - Product Information

Application FC, WB,E
Primary Accession Q9UK99

Other Accession <u>D4ABP9</u>, <u>Q9DC63</u>, <u>A6H7H7</u>

Reactivity Human

Predicted Bovine, Mouse, Rat

Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Calculated MW 54561
Antigen Region 395-422

## FBXO3 Antibody (C-term) - Additional Information

**Gene ID 26273** 

#### **Other Names**

F-box only protein 3, FBXO3, FBX3

#### Target/Specificity

This FBXO3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 395-422 amino acids from the C-terminal region of human FBXO3.

#### **Dilution**

FC~~1:10~50 WB~~1:1000

E~~Use at an assay dependent concentration.

#### **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### **Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### **Precautions**

FBXO3 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

#### FBXO3 Antibody (C-term) - Protein Information

Name FBXO3 (HGNC:13582)





### Synonyms FBX3

**Function** Substrate recognition component of the SCF (SKP1-CUL1-F-box protein)-type E3 ubiquitin ligase complex, SCF(FBXO3), which mediates the ubiquitination and subsequent proteasomal degradation of target proteins (PubMed:18809579, PubMed:26037928). Mediates the ubiquitination of HIPK2 and probably that of EP300, leading to rapid degradation by the proteasome (PubMed:18809579). In the presence of PML, HIPK2 ubiquitination still occurs, but degradation is prevented (PubMed:18809579). PML, HIPK2 and FBXO3 may act synergically to activate p53/TP53-dependent transactivation (PubMed:18809579). The SCF(FBXO3) also acts as a regulator of inflammation by mediating ubiquitination and degradation of FBXL2 in response to lipopolysaccharide (LPS) (PubMed:26037928, PubMed:27010866). The SCF(FBXO3) complex specifically recognizes FBXL2 phosphorylated at 'Thr-404' and promotes its ubiquitination (PubMed:27010866).

#### **Cellular Location**


Nucleus. Note=Colocalizes with PML at the peripheries of nuclear bodies

#### FBXO3 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

# FBXO3 Antibody (C-term) - Images



Western blot analysis of FBXO3 Antibody (C-term) (Cat. #AP9195b) in HepG2 cell line lysates (35ug/lane). FBXO3 (arrow) was detected using the purified Pab.





FBXO3 Antibody (C-term) (Cat. #AP9195b) flow cytometric analysis of HepG2 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

## FBXO3 Antibody (C-term) - Background

FBXO3 encodes a member of the F-box protein family which is characterized by an approximately 40 amino acid motif, the F-box. The F-box proteins constitute one of the four subunits of the ubiquitin protein ligase complex called SCFs (SKP1-cullin-F-box), which function in phosphorylation-dependent ubiquitination. The F-box proteins are divided into 3 classes: Fbws containing WD-40 domains, Fbls containing leucine-rich repeats, and Fbxs containing either different protein-protein interaction modules or no recognizable motifs.

## FBXO3 Antibody (C-term) - References

Shima,Y., et.al., Mol. Cell. Biol. 28 (23), 7126-7138 (2008) llyin,G.P., et.al., Genomics 67 (1), 40-47 (2000)

FBXO3 Antibody (C-term) - Citations

• Targeted inhibition of the COP9 signalosome for treatment of cancer.