

PERP Antibody
Rabbit mAb
Catalog # AP91994**Specification**

PERP Antibody - Product Information

Application	WB, ICC
Primary Accession	Q96FX8
Reactivity	Rat
Clonality	Monoclonal
Other Names	
KCP1; KRTCAP1; Perp; PIGPC1; THW;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	21386 Da

PERP Antibody - Additional Information

Dilution	WB~~1:1000 ICC~~N/A
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human PERP
Description	Component of intercellular desmosome junctions. Plays a role in stratified epithelial integrity and cell-cell adhesion by promoting desmosome assembly. Plays a role as an effector in the TP53-dependent apoptotic pathway.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

PERP Antibody - Protein Information**Name** PERP ([HGNC:17637](#))**Function**

Component of intercellular desmosome junctions (By similarity). Plays a role in stratified epithelial integrity and cell- cell adhesion by promoting desmosome assembly (By similarity). Thereby plays a role in barrier function of the skin against infection (By similarity). Plays a role in mammary epithelial tissue homeostasis and remodeling during and after pregnancy, potentially via its involvement in desmosome cell-cell junctions (By similarity). Required for tooth enamel development via facilitating desmosome-mediated ameloblast adhesion to the stratum intermedium during the transitional stage of amelogenesis (By similarity). May also play a role in downstream transcriptional regulation of other genes involved in amelogenesis such as AMBN,

ENAM, MMP20 and KLK4 (By similarity). Plays a role as an effector in the TP53-dependent apoptotic pathway (By similarity). Positively regulates apoptosis in T-helper 17 (Th17) cell populations via caspase-dependent signaling (By similarity). Promotes neutrophil transepithelial migration in response to chemoattractants such as heparin A3 (HXA3), N-Formylmethionyl-leucyl-phenylalanine (fMLP) and CXCL8/IL-8 (PubMed:25486861). Required for neutrophil transepithelial migration in response to S.typhimurium infection (PubMed:25486861). May act as a positive regulator of endothelial cell apoptosis in response to blood flow-derived shear stress (By similarity).

Cellular Location

Cell junction, desmosome {ECO:0000250|UniProtKB:Q9JK95}. Cell membrane; Multi-pass membrane protein. Cytoplasm. Note=Associated with desmosomes (By similarity). Colocalizes with KRT14 in the cell membrane (PubMed:31898316). Clusters in a punctate pattern throughout the epithelial cytoplasm, in response to S.typhimurium infection (PubMed:25486861). {ECO:0000250|UniProtKB:Q9JK95, ECO:0000269|PubMed:25486861, ECO:0000269|PubMed:31898316}

Tissue Location

Expressed in skin, heart, placental, liver, pancreas, keratinocytes and dermal fibroblasts. May translocate to the intestinal apical epithelial cell surface via sipA and sctB1/sipC- promoted exocytic translocation following infection by S. Typhimurium (PubMed:25486861, PubMed:27078059).


PERP Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PERP Antibody - Images





Western blot analysis of PERP expression in A431 cell lysate.