

Cytochrome P450 4A Antibody
Rabbit mAb
Catalog # AP92005**Specification****Cytochrome P450 4A Antibody - Product Information**

Application	WB, IHC, IP
Primary Accession	Q02928
Reactivity	Rat
Clonality	Monoclonal
Other Names	
CYP4A; Cyp4a1; Cyp4a10; CYP4A11; Cyp4a14; Cyp4a3; CYP4A7; CYP4AII; CYPIVA11;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	59348 Da

Cytochrome P450 4A Antibody - Additional Information

Dilution	WB~~1:1000 IHC~~1:100~500 IP~~N/A
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Cytochrome P450 4A
Description	Catalyzes the omega- and (omega-1)-hydroxylation of various fatty acids such as laurate, myristate and palmitate. Has little activity toward prostaglandins A1 and E1. Oxidizes arachidonic acid to 20-hydroxyeicosatetraenoic acid (20-HETE).
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Cytochrome P450 4A Antibody - Protein Information**Name** CYP4A11 {ECO:0000303|PubMed:8274222, ECO:0000312|HGNC:HGNC:2642}**Function**

A cytochrome P450 monooxygenase involved in the metabolism of fatty acids and their oxygenated derivatives (oxylipins) (PubMed:10553002, PubMed:10660572, PubMed:15611369, PubMed:1739747)

target="_blank">>1739747, PubMed:>7679927, PubMed:>8914854). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (CPR; NADPH-ferredoxin reductase) (PubMed:>10553002, PubMed:>10660572, PubMed:>15611369, PubMed:>1739747, PubMed:>7679927, PubMed:>8914854). Catalyzes predominantly the oxidation of the terminal carbon (omega-oxidation) of saturated and unsaturated fatty acids, the catalytic efficiency decreasing in the following order: dodecanoic > tetradecanoic > (9Z)-octadecenoic > (9Z,12Z)-octadecadienoic > hexadecanoic acid (PubMed:>10553002, PubMed:>10660572). Acts as a major omega-hydroxylase for dodecanoic (lauric) acid in liver (PubMed:>15611369, PubMed:>1739747, PubMed:>7679927, PubMed:>8914854). Participates in omega-hydroxylation of (5Z,8Z,11Z,14Z)-eicosatetraenoic acid (arachidonate) to 20-hydroxyeicosatetraenoic acid (20-HETE), a signaling molecule acting both as vasoconstrictive and natriuretic with overall effect on arterial blood pressure (PubMed:>10620324, PubMed:>10660572, PubMed:>15611369). Can also catalyze the oxidation of the penultimate carbon (omega-1 oxidation) of fatty acids with lower efficiency (PubMed:>7679927). May contribute to the degradation of saturated very long-chain fatty acids (VLCFAs) such as docosanoic acid, by catalyzing successive omega-oxidations to the corresponding dicarboxylic acid, thereby initiating chain shortening (PubMed:>18182499). Omega-hydroxylates (9R,10S)-epoxy-octadecanoate stereoisomer (PubMed:>15145985). Plays a minor role in omega-oxidation of long-chain 3-hydroxy fatty acids (PubMed:>18065749). Has little activity toward prostaglandins A1 and E1 (PubMed:>7679927).

Cellular Location

Endoplasmic reticulum membrane; Peripheral membrane protein. Microsome membrane; Peripheral membrane protein

Tissue Location


Expressed in liver (PubMed:7679927). Expressed in S2 and S3 segments of proximal tubules in cortex and outer medulla of kidney (PubMed:10660572, PubMed:7679927).

Cytochrome P450 4A Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Cytochrome P450 4A Antibody - Images

Western blot analysis of Cytochrome P450 4A expression in Human fetal kidney lysate.