

**CD53 Antibody**  
**Rabbit mAb**  
**Catalog # AP92058****Specification****CD53 Antibody - Product Information**

Application	WB, IHC, FC, ICC
Primary Accession	<a href="#">P19397</a>
Clonality	Monoclonal
<b>Other Names</b>	
CD53; CD53 antigen; MOX44; OX44; RATOX44; Tetraspanin25; Transmembrane glycoprotein; TSPAN25; PRO1340;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	24341 Da

**CD53 Antibody - Additional Information**

Dilution	WB~~1:1000 IHC~~1:100~500 FC~~1:10~50 ICC~~N/A
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human CD53
Description	May be involved in growth regulation in hematopoietic cells.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

**CD53 Antibody - Protein Information****Name** CD53**Synonyms** MOX44, TSPAN25**Function**

Structural component of specialized membrane microdomains known as tetraspanin-enriched microdomains (TERMs), which act as platforms for receptor clustering and signaling (PubMed:<a href="http://www.uniprot.org/citations/28487417" target="\_blank">28487417</a>). Participates thereby in diverse biological functions such as cell signal transduction, adhesion, migration and protein trafficking (PubMed:<a href="http://www.uniprot.org/citations/32974937" target="\_blank">32974937</a>, PubMed:<a href="http://www.uniprot.org/citations/35767951" target="\_blank">35767951</a>). Plays a role in the activation of monocytes and B-cells (PubMed:<a href="http://www.uniprot.org/citations/8335905" target="\_blank">8335905</a>).

Acts as an essential regulator of B-cell development by promoting interleukin-7 receptor/IL7R signaling (By similarity). Also promotes, in B-cells, the BCR signaling by recruiting PKC to the plasma membrane in order to phosphorylate its substrates (PubMed:<a href="http://www.uniprot.org/citations/28487417" target="\_blank">28487417</a>). Plays an essential role in B- and T-cells homing to lymph nodes by stabilizing L-selectin/SELL cell surface expression (By similarity). Also mediates metabolic and inflammatory functions in hepatocytes and adipose tissue by promoting TNF-alpha and LPS signaling independent of the immune compartment (By similarity).

#### Cellular Location

Cell membrane. Cell junction {ECO:0000250|UniProtKB:Q61451}. Membrane; Multi-pass membrane protein. Synapse. Note=Concentrates in localized microdomains along the plasma membrane at the contact sites between cells of fused myotubes.  
{ECO:0000250|UniProtKB:Q61451}

#### Tissue Location


B-cells, monocytes, macrophages, neutrophils, single (CD4 or CD8) positive thymocytes and peripheral T-cells

### CD53 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### CD53 Antibody - Images



Western blot analysis of CD53 expression in Jurkat cell lysate.