

**TFE3 Antibody**  
**Rabbit mAb**  
**Catalog # AP92080****Specification**

---

**TFE3 Antibody - Product Information**

Application	IHC, FC, ICC
Primary Accession	<a href="#">P19532</a>
Clonality	Monoclonal
<b>Other Names</b>	
bHLHe33; RCCP2; RCCX1; Tcfe3; Tfe3; TFEA;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	61521 Da

**TFE3 Antibody - Additional Information**

Dilution	IHC~~1:100~500 FC~~1:10~50 ICC~~N/A
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human TFE3
Description	Transcription factor that specifically recognizes and binds E-box sequences (3'-CANNTG-5'). Efficient DNA-binding requires dimerization with itself or with another MiT/TFE family member such as TFEB or MITF. In association with TFEB, activates the expression of CD40L in T-cells, thereby playing a role in T-cell-dependent antibody responses in activated CD4(+) T-cells and thymus-dependent humoral immunity.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

**TFE3 Antibody - Protein Information**

**Name** TFE3 {ECO:0000303|PubMed:9393982, ECO:0000312|HGNC:HGNC:11752}

**Function**

Transcription factor that acts as a master regulator of lysosomal biogenesis and immune response (PubMed:<a href="http://www.uniprot.org/citations/2338243" target="\_blank">2338243</a>, PubMed:<a href="http://www.uniprot.org/citations/24448649" target="\_blank">24448649</a>),

PubMed:<a href="http://www.uniprot.org/citations/29146937" target="\_blank">29146937</a>, PubMed:<a href="http://www.uniprot.org/citations/30733432" target="\_blank">30733432</a>, PubMed:<a href="http://www.uniprot.org/citations/31672913" target="\_blank">31672913</a>, PubMed:<a href="http://www.uniprot.org/citations/37079666" target="\_blank">37079666</a>). Specifically recognizes and binds E-box sequences (5'-CANNTG-3'); efficient DNA-binding requires dimerization with itself or with another MiT/TFE family member such as TFEB or MITF (PubMed:<a href="http://www.uniprot.org/citations/24448649" target="\_blank">24448649</a>). Involved in the cellular response to amino acid availability by acting downstream of MTOR: in the presence of nutrients, TFE3 phosphorylation by MTOR promotes its inactivation (PubMed:<a href="http://www.uniprot.org/citations/24448649" target="\_blank">24448649</a>, PubMed:<a href="http://www.uniprot.org/citations/31672913" target="\_blank">31672913</a>, PubMed:<a href="http://www.uniprot.org/citations/36608670" target="\_blank">36608670</a>). Upon starvation or lysosomal stress, inhibition of MTOR induces TFE3 dephosphorylation, resulting in transcription factor activity (PubMed:<a href="http://www.uniprot.org/citations/24448649" target="\_blank">24448649</a>, PubMed:<a href="http://www.uniprot.org/citations/31672913" target="\_blank">31672913</a>, PubMed:<a href="http://www.uniprot.org/citations/36608670" target="\_blank">36608670</a>). Specifically recognizes and binds the CLEAR-box sequence (5'-GTCACGTGAC-3') present in the regulatory region of many lysosomal genes, leading to activate their expression, thereby playing a central role in expression of lysosomal genes (PubMed:<a href="http://www.uniprot.org/citations/24448649" target="\_blank">24448649</a>). Maintains the pluripotent state of embryonic stem cells by promoting the expression of genes such as ESRRB; mTOR- dependent TFE3 cytosolic retention and inactivation promotes exit from pluripotency (By similarity). Required to maintain the naive pluripotent state of hematopoietic stem cell; mTOR-dependent cytoplasmic retention of TFE3 promotes the exit of hematopoietic stem cell from pluripotency (PubMed:<a href="http://www.uniprot.org/citations/30733432" target="\_blank">30733432</a>). TFE3 activity is also involved in the inhibition of neuronal progenitor differentiation (By similarity). Acts as a positive regulator of browning of adipose tissue by promoting expression of target genes; mTOR-dependent phosphorylation promotes cytoplasmic retention of TFE3 and inhibits browning of adipose tissue (By similarity). In association with TFEB, activates the expression of CD40L in T-cells, thereby playing a role in T-cell- dependent antibody responses in activated CD4(+) T-cells and thymus- dependent humoral immunity (By similarity). Specifically recognizes the MUE3 box, a subset of E-boxes, present in the immunoglobulin enhancer (PubMed:<a href="http://www.uniprot.org/citations/2338243" target="\_blank">2338243</a>). It also binds very well to a USF/MLTF site (PubMed:<a href="http://www.uniprot.org/citations/2338243" target="\_blank">2338243</a>). Promotes TGF-beta-induced transcription of COL1A2; via its interaction with TSC22D1 at E-boxes in the gene proximal promoter (By similarity). May regulate lysosomal positioning in response to nutrient deprivation by promoting the expression of PIP4P1 (PubMed:<a href="http://www.uniprot.org/citations/29146937" target="\_blank">29146937</a>).

### Cellular Location

Cytoplasm, cytosol. Nucleus. Lysosome membrane. Note=When nutrients are present, recruited to the lysosomal membrane via association with GDP-bound RagC/RRAGC (or RagD/RRAGD): it is then phosphorylated by MTOR (PubMed:24448649, PubMed:37079666). Phosphorylation by MTOR prevents nuclear translocation and promotes ubiquitination and degradation (PubMed:22692423, PubMed:30733432, PubMed:36608670, PubMed:37079666) Conversely, inhibition of mTORC1, starvation and lysosomal disruption, promotes dephosphorylation and translocation to the nucleus (PubMed:22692423, PubMed:30733432, PubMed:37079666)

### Tissue Location

Ubiquitous in fetal and adult tissues.

### TFE3 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **TFE3 Antibody - Images**