

**METTL3 Antibody**  
**Rabbit mAb**  
**Catalog # AP92110****Specification**

---

**METTL3 Antibody - Product Information**

Application	WB, IHC, FC, ICC, IP
Primary Accession	<a href="#">Q86U44</a>
Reactivity	Rat
Clonality	Monoclonal
<b>Other Names</b>	
IME4; M6A; Methyltransferase like protein 3; METTL3; MTA70; N6 adenosine methyltransferase 70 kDa subunit;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	64474 Da

**METTL3 Antibody - Additional Information**

Dilution	WB~~1:1000 IHC~~1:100~500 FC~~1:10~50 ICC~~N/A IP~~N/A
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human METTL3
Description	N6-methyltransferase that methylates adenosine residues of some mRNAs. N6-methyladenosine (m6A), which is present at internal sites of some mRNAs, may play a role in the efficiency of mRNA splicing, transport or translation.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

**METTL3 Antibody - Protein Information****Name** METTL3 ([HGNC:17563](#))**Synonyms** MTA70**Function**

The METTL3-METTL14 heterodimer forms a N6-methyltransferase complex that methylates adenosine residues at the N(6) position of some RNAs and regulates various processes such as the

circadian clock, differentiation of embryonic and hematopoietic stem cells, cortical neurogenesis, response to DNA damage, differentiation of T-cells and primary miRNA processing (PubMed:<a href="http://www.uniprot.org/citations/22575960" target="\_blank">22575960</a>, PubMed:<a href="http://www.uniprot.org/citations/24284625" target="\_blank">24284625</a>, PubMed:<a href="http://www.uniprot.org/citations/25719671" target="\_blank">25719671</a>, PubMed:<a href="http://www.uniprot.org/citations/25799998" target="\_blank">25799998</a>, PubMed:<a href="http://www.uniprot.org/citations/26321680" target="\_blank">26321680</a>, PubMed:<a href="http://www.uniprot.org/citations/26593424" target="\_blank">26593424</a>, PubMed:<a href="http://www.uniprot.org/citations/27281194" target="\_blank">27281194</a>, PubMed:<a href="http://www.uniprot.org/citations/27373337" target="\_blank">27373337</a>, PubMed:<a href="http://www.uniprot.org/citations/27627798" target="\_blank">27627798</a>, PubMed:<a href="http://www.uniprot.org/citations/28297716" target="\_blank">28297716</a>, PubMed:<a href="http://www.uniprot.org/citations/29348140" target="\_blank">29348140</a>, PubMed:<a href="http://www.uniprot.org/citations/29506078" target="\_blank">29506078</a>, PubMed:<a href="http://www.uniprot.org/citations/30428350" target="\_blank">30428350</a>, PubMed:<a href="http://www.uniprot.org/citations/9409616" target="\_blank">9409616</a>). In the heterodimer formed with METTL14, METTL3 constitutes the catalytic core (PubMed:<a href="http://www.uniprot.org/citations/27281194" target="\_blank">27281194</a>, PubMed:<a href="http://www.uniprot.org/citations/27373337" target="\_blank">27373337</a>, PubMed:<a href="http://www.uniprot.org/citations/27627798" target="\_blank">27627798</a>). N6-methyladenosine (m6A), which takes place at the 5'-[AG]GAC-3' consensus sites of some mRNAs, plays a role in mRNA stability, processing, translation efficiency and editing (PubMed:<a href="http://www.uniprot.org/citations/22575960" target="\_blank">22575960</a>, PubMed:<a href="http://www.uniprot.org/citations/24284625" target="\_blank">24284625</a>, PubMed:<a href="http://www.uniprot.org/citations/25719671" target="\_blank">25719671</a>, PubMed:<a href="http://www.uniprot.org/citations/25799998" target="\_blank">25799998</a>, PubMed:<a href="http://www.uniprot.org/citations/26321680" target="\_blank">26321680</a>, PubMed:<a href="http://www.uniprot.org/citations/26593424" target="\_blank">26593424</a>, PubMed:<a href="http://www.uniprot.org/citations/28297716" target="\_blank">28297716</a>, PubMed:<a href="http://www.uniprot.org/citations/9409616" target="\_blank">9409616</a>). M6A acts as a key regulator of mRNA stability: methylation is completed upon the release of mRNA into the nucleoplasm and promotes mRNA destabilization and degradation (PubMed:<a href="http://www.uniprot.org/citations/28637692" target="\_blank">28637692</a>). In embryonic stem cells (ESCs), m6A methylation of mRNAs encoding key naive pluripotency-promoting transcripts results in transcript destabilization, promoting differentiation of ESCs (By similarity). M6A regulates the length of the circadian clock: acts as an early pace-setter in the circadian loop by putting mRNA production on a fast-track for facilitating nuclear processing, thereby providing an early point of control in setting the dynamics of the feedback loop (By similarity). M6A also regulates circadian regulation of hepatic lipid metabolism (PubMed:<a href="http://www.uniprot.org/citations/30428350" target="\_blank">30428350</a>). M6A regulates spermatogonial differentiation and meiosis and is essential for male fertility and spermatogenesis (By similarity). Also required for oogenesis (By similarity). Involved in the response to DNA damage: in response to ultraviolet irradiation, METTL3 rapidly catalyzes the formation of m6A on poly(A) transcripts at DNA damage sites, leading to the recruitment of POLK to DNA damage sites (PubMed:<a href="http://www.uniprot.org/citations/28297716" target="\_blank">28297716</a>). M6A is also required for T-cell homeostasis and differentiation: m6A methylation of transcripts of SOCS family members (SOCS1, SOCS3 and CISH) in naive T-cells promotes mRNA destabilization and degradation, promoting T-cell differentiation (By similarity). Inhibits the type I interferon response by mediating m6A methylation of IFNB (PubMed:<a href="http://www.uniprot.org/citations/30559377" target="\_blank">30559377</a>). M6A also takes place in other RNA molecules, such as primary miRNA (pri- miRNAs) (PubMed:<a href="http://www.uniprot.org/citations/25799998" target="\_blank">25799998</a>). Mediates m6A methylation of Xist RNA, thereby participating in random X inactivation: m6A methylation of Xist leads to target YTHDC1 reader on Xist and promote transcription repression activity of Xist (PubMed:<a href="http://www.uniprot.org/citations/27602518" target="\_blank">27602518</a>). M6A also regulates cortical neurogenesis: m6A methylation of transcripts related to transcription factors, neural stem cells, the cell cycle and neuronal differentiation during brain development

promotes their destabilization and decay, promoting differentiation of radial glial cells (By similarity). METTL3 mediates methylation of pri-miRNAs, marking them for recognition and processing by DGCR8 (PubMed:<a href="http://www.uniprot.org/citations/25799998" target="\_blank">25799998</a>). Acts as a positive regulator of mRNA translation independently of the methyltransferase activity: promotes translation by interacting with the translation initiation machinery in the cytoplasm (PubMed:<a href="http://www.uniprot.org/citations/27117702" target="\_blank">27117702</a>). Its overexpression in a number of cancer cells suggests that it may participate in cancer cell proliferation by promoting mRNA translation (PubMed:<a href="http://www.uniprot.org/citations/27117702" target="\_blank">27117702</a>). During human coronavirus SARS-CoV-2 infection, adds m6A modifications in SARS-CoV-2 RNA leading to decreased RIGI binding and subsequently dampening the sensing and activation of innate immune responses (PubMed:<a href="http://www.uniprot.org/citations/33961823" target="\_blank">33961823</a>).

#### **Cellular Location**

Nucleus. Nucleus speckle. Cytoplasm. Note=Colocalizes with speckles in interphase nuclei, suggesting that it may be associated with nuclear pre-mRNA splicing components (PubMed:9409616). In response to ultraviolet irradiation, colocalizes to DNA damage sites however, it probably does not bind DNA but localizes in the vicinity of DNA damage sites (PubMed:28297716).

#### **Tissue Location**


Widely expressed at low level. Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon and peripheral blood leukocytes.


#### **METTL3 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **METTL3 Antibody - Images**





Western blot analysis of METTL3 expression in (1) 293 cell lysate; (2) RAW 264.7 cell lysate.