

CaMKI Antibody

Rabbit mAb Catalog # AP92130

Specification

CaMKI Antibody - Product Information

Application WB, IHC, ICC **Primary Accession** 014012 Reactivity Rat Clonality Monoclonal **Other Names** CaM K1; CaM KI; CaMK 1; CAMK I; CaMK1 alpha; CAMK1 PEN; CaMKI alpha;

Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	41337 Da

CaMKI Antibody - Additional Information

Dilution	WB~~1:1000
	IHC~~1:100~500
	ICC~~N/A
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human
	CaMKI
Description	Calcium/calmodulin-dependent protein
	kinase that operates in the
	calcium-triggered CaMKK-CaMK1 signaling
	cascade and, upon calcium influx,
	regulates transcription activators activity,
	cell cycle, hormone production, cell
	differentiation, actin filament organization
	and neurite outgrowth. Recognizes the
	substrate consensus sequence
	[MVLIF]-x-R-x(2)-[ST]-x(3)-[MVLIF].
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline ,
	pH 7.4, 150mM NaCl, 0.02% sodium azide
	and 50% glycerol. Store at +4°C short
	term. Store at -20°C long term. Avoid
	freeze / thaw cycle.

CaMKI Antibody - Protein Information

Name CAMK1

Function

Calcium/calmodulin-dependent protein kinase that operates in the calcium-triggered CaMKK-CaMK1 signaling cascade and, upon calcium influx, regulates transcription activators activity, cell cycle, hormone production, cell differentiation, actin filament organization and neurite



outgrowth. Recognizes the substrate consensus sequence [MVLIF]-x-R-x(2)-[ST]-x(3)-[MVLIF]. Regulates axonal extension and growth cone motility in hippocampal and cerebellar nerve cells. Upon NMDA receptor-mediated Ca(2+) elevation, promotes dendritic growth in hippocampal neurons and is essential in synapses for full long-term potentiation (LTP) and ERK2-dependent translational activation. Downstream of NMDA receptors, promotes the formation of spines and synapses in hippocampal neurons by phosphorylating ARHGEF7/BETAPIX on 'Ser-694', which results in the enhancement of ARHGEF7 activity and activation of RAC1. Promotes neuronal differentiation and neurite outgrowth by activation and phosphorylation of MARK2 on 'Ser-91', 'Ser-92', 'Ser-93' and 'Ser-294'. Promotes nuclear export of HDAC5 and binding to 14-3-3 by phosphorylation of 'Ser-259' and 'Ser-498' in the regulation of muscle cell differentiation. Regulates NUMB-mediated endocytosis by phosphorylation of NUMB on 'Ser-276' and 'Ser-295'. Involved in the regulation of basal and estrogen-stimulated migration of medulloblastoma cells through ARHGEF7/BETAPIX phosphorylation (By similarity). Is required for proper activation of cvclin-D1/CDK4 complex during G1 progression in diploid fibroblasts. Plays a role in K(+) and ANG2-mediated regulation of the aldosterone synthase (CYP11B2) to produce aldosterone in the adrenal cortex. Phosphorylates EIF4G3/eIF4GII. In vitro phosphorylates CREB1, ATF1, CFTR, MYL9 and SYN1/synapsin I.

Cellular Location Cytoplasm. Nucleus. Note=Predominantly cytoplasmic.

Tissue Location Widely expressed. Expressed in cells of the zona glomerulosa of the adrenal cortex.

CaMKI Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

CaMKI Antibody - Images





Western blot analysis of CaMKI expression in SH-SY5Y cell lysate.