

LRP6 Antibody
Rabbit mAb
Catalog # AP92131**Specification****LRP6 Antibody - Product Information**

Application	WB, FC, ICC
Primary Accession	075581
Reactivity	Rat
Clonality	Monoclonal
Other Names	
LRP6;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	180429 Da

LRP6 Antibody - Additional Information

Dilution	WB~~1:1000 FC~~1:10~50 ICC~~N/A
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human LRP6
Description	Component of the Wnt-Fzd-LRP5-LRP6 complex that triggers beta-catenin signaling through inducing aggregation of receptor-ligand complexes into ribosome-sized signalosomes. Cell-surface coreceptor of Wnt/beta-catenin signaling, which plays a pivotal role in bone formation.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

LRP6 Antibody - Protein Information**Name** LRP6**Function**

Component of the Wnt-Fzd-LRP5-LRP6 complex that triggers beta-catenin signaling through inducing aggregation of receptor-ligand complexes into ribosome-sized signalosomes (PubMed:<a href="<http://www.uniprot.org/citations/11357136>">11357136, PubMed:<a href="<http://www.uniprot.org/citations/11448771>">11448771, PubMed:<a href="<http://www.uniprot.org/citations/15778503>">15778503, PubMed:[15778503](#)

href="http://www.uniprot.org/citations/16341017" target="_blank">>16341017, PubMed:>16513652, PubMed:>17326769, PubMed:>17400545, PubMed:>19107203, PubMed:>19293931, PubMed:>19801552, PubMed:>28341812). Cell-surface coreceptor of Wnt/beta- catenin signaling, which plays a pivotal role in bone formation (PubMed:>11357136, PubMed:>11448771, PubMed:>15778503, PubMed:>16341017, PubMed:>16513652, PubMed:>17326769, PubMed:>17400545, PubMed:>19107203, PubMed:>19293931, PubMed:>19801552, PubMed:>28341812). The Wnt-induced Fzd/LRP6 coreceptor complex recruits DVL1 polymers to the plasma membrane which, in turn, recruits the AXIN1/GSK3B-complex to the cell surface promoting the formation of signalosomes and inhibiting AXIN1/GSK3-mediated phosphorylation and destruction of beta-catenin (PubMed:>16513652). Required for posterior patterning of the epiblast during gastrulation (By similarity).

Cellular Location

Cell membrane; Single-pass type I membrane protein Endoplasmic reticulum. Membrane raft. Note=On Wnt signaling, undergoes a cycle of caveolin- or clathrin-mediated endocytosis and plasma membrane location. Released from the endoplasmic reticulum on palmitoylation Mono-ubiquitination retains it in the endoplasmic reticulum in the absence of palmitoylation. On Wnt signaling, phosphorylated, aggregates and colocalizes with AXIN1 and GSK3B at the plasma membrane in LRP6- signalosomes (By similarity). Chaperoned to the plasma membrane by HSP90B1 and MESD (PubMed:23572575). {ECO:0000250|UniProtKB:O88572, ECO:0000269|PubMed:23572575}

Tissue Location


Widely coexpressed with LRP5 during embryogenesis and in adult tissues

LRP6 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

LRP6 Antibody - Images



Western blot analysis of LRP6 expression in Jurkat cell lysate.