

GluA4 Antibody
Rabbit mAb
Catalog # AP92132**Specification**

GluA4 Antibody - Product Information

| | |
|--|------------------------|
| Application | WB, FC |
| Primary Accession | P48058 |
| Reactivity | Rat |
| Clonality | Monoclonal |
| Other Names | |
| AMPA4; GluA4; GluR4; GLUR4C; GLURD; Gria4; | |
| Isotype | Rabbit IgG |
| Host | Rabbit |
| Calculated MW | 100871 Da |

GluA4 Antibody - Additional Information

| | |
|------------------------------|--|
| Dilution | WB~~1:1000 FC~~1:10~50 |
| Purification | Affinity-chromatography |
| Immunogen | A synthesized peptide derived from human GluA4 |
| Description | Ionotropic glutamate receptor. L-glutamate acts as an excitatory neurotransmitter at many synapses in the central nervous system. Binding of the excitatory neurotransmitter L-glutamate induces a conformation change, leading to the opening of the cation channel, and thereby converts the chemical signal to an electrical impulse. |
| Storage Condition and Buffer | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle. |

GluA4 Antibody - Protein Information

Name GRIA4 {ECO:0000303|PubMed:29220673, ECO:0000312|HGNC:HGNC:4574}

Function

Ionotropic glutamate receptor that functions as a ligand- gated cation channel, gated by L-glutamate and glutamatergic agonists such as alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA), quisqualic acid, and kainic acid (By similarity). L-glutamate acts as an excitatory neurotransmitter at many synapses in the central nervous system and plays an important role in fast excitatory synaptic transmission (By similarity).

Binding of the excitatory neurotransmitter L-glutamate induces a conformation change, leading to the opening of the cation channel, and thereby converts the chemical signal to an electrical impulse upon entry of monovalent and divalent cations such as sodium and calcium. The receptor then desensitizes rapidly and enters a transient inactive state, characterized by the presence of bound agonist (By similarity). In the presence of CACNG8, shows resensitization which is characterized by a delayed accumulation of current flux upon continued application of L-glutamate (PubMed: [21172611](http://www.uniprot.org/citations/21172611)).

Cellular Location


Cell membrane {ECO:0000250|UniProtKB:P19493}; Multi-pass membrane protein {ECO:0000250|UniProtKB:P19493} Postsynaptic cell membrane {ECO:0000250|UniProtKB:P19493}; Multi-pass membrane protein {ECO:0000250|UniProtKB:P19493}. Cell projection, dendrite {ECO:0000250|UniProtKB:P19493}. Postsynaptic cell membrane {ECO:0000250|UniProtKB:P42262}; Multi-pass membrane protein {ECO:0000250|UniProtKB:P42262}

GluA4 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

GluA4 Antibody - Images



Western blot analysis of Ionotropic Glutamate receptor 4 expression in SH-SY5Y cell lysate.