

Lamin A/C Antibody
Rabbit mAb
Catalog # AP92257**Specification****Lamin A/C Antibody - Product Information**

Application	WB, IHC, FC, ICC
Primary Accession	P02545
Reactivity	Rat
Clonality	Monoclonal
Other Names	
CDCD1; CDDC; CMD1A; CMT2B1; EMD2; FPL; FPLD; FPLD2; HGPS; IDC; Lamin A; Lamin A/C; Lamin C; LDP1; LFP; LGMD1B; LMN1; LMNA; LMNC; LMNL1; PRO1;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	74139 Da

Lamin A/C Antibody - Additional Information

Dilution	WB~~~1:1000 IHC~~~1:100~500 FC~~~1:10~50 ICC~~~N/A
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Lamin A/C
Description	Lamins are components of the nuclear lamina, a fibrous layer on the nucleoplasmic side of the inner nuclear membrane, which is thought to provide a framework for the nuclear envelope and may also interact with chromatin.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Lamin A/C Antibody - Protein Information**Name** LMNA**Synonyms** LMN1**Function**

[Lamin-A/C]: Lamins are intermediate filament proteins that assemble into a filamentous meshwork, and which constitute the major components of the nuclear lamina, a fibrous layer on the nucleoplasmic side of the inner nuclear membrane (PubMed:>10080180, PubMed:>10580070, PubMed:>10587585, PubMed:>10814726, PubMed:>11799477, PubMed:>12075506, PubMed:>12927431, PubMed:>15317753, PubMed:>18551513, PubMed:>18611980, PubMed:>2188730, PubMed:>22431096, PubMed:>2344612, PubMed:>23666920, PubMed:>24741066, PubMed:>31434876, PubMed:>31548606, PubMed:>37788673, PubMed:>37832547). Lamins provide a framework for the nuclear envelope, bridging the nuclear envelope and chromatin, thereby playing an important role in nuclear assembly, chromatin organization, nuclear membrane and telomere dynamics (PubMed:>10080180, PubMed:>10580070, PubMed:>10587585, PubMed:>10814726, PubMed:>11799477, PubMed:>12075506, PubMed:>12927431, PubMed:>15317753, PubMed:>18551513, PubMed:>18611980, PubMed:>22431096, PubMed:>23666920, PubMed:>24741066, PubMed:>31548606, PubMed:>37788673, PubMed:>37832547). Lamin A and C also regulate matrix stiffness by conferring nuclear mechanical properties (PubMed:>23990565, PubMed:>25127216). The structural integrity of the lamina is strictly controlled by the cell cycle, as seen by the disintegration and formation of the nuclear envelope in prophase and telophase, respectively (PubMed:>2188730, PubMed:>2344612). Lamin A and C are present in equal amounts in the lamina of mammals (PubMed:>10080180, PubMed:>10580070, PubMed:>10587585, PubMed:>10814726, PubMed:>11799477, PubMed:>12075506, PubMed:>12927431, PubMed:>15317753, PubMed:>18551513, PubMed:>18611980, PubMed:>22431096, PubMed:>23666920

target="_blank">>23666920, PubMed:31548606). Also involved in DNA repair: recruited by DNA repair proteins XRCC4 and IFFO1 to the DNA double-strand breaks (DSBs) to prevent chromosome translocation by immobilizing broken DNA ends (PubMed:31548606). Required for normal development of peripheral nervous system and skeletal muscle and for muscle satellite cell proliferation (PubMed:10080180, PubMed:10814726, PubMed:11799477, PubMed:18551513, PubMed:22431096). Required for osteoblastogenesis and bone formation (PubMed:12075506, PubMed:15317753, PubMed:18611980). Also prevents fat infiltration of muscle and bone marrow, helping to maintain the volume and strength of skeletal muscle and bone (PubMed:10587585). Required for cardiac homeostasis (PubMed:10580070, PubMed:12927431, PubMed:18611980, PubMed:23666920).

Cellular Location

Nucleus lamina. Nucleus envelope. Nucleus, nucleoplasm. Nucleus matrix. Note=Farnesylation of prelamin-A/C facilitates nuclear envelope targeting and subsequent cleavage by ZMPSTE24/FACE1 to remove the farnesyl group produces mature lamin-A/C, which can then be inserted into the nuclear lamina (PubMed:15317753) EMD is required for proper localization of non-farnesylated prelamin- A/C (PubMed:19323649). Also localizes to the micronuclear envelope in response to response to genome instability (PubMed:37788673)

Tissue Location


In the arteries, prelamin-A/C accumulation is not observed in young healthy vessels but is prevalent in medial vascular smooth muscle cells (VSMCs) from aged individuals and in atherosclerotic lesions, where it often colocalizes with senescent and degenerate VSMCs. Prelamin-A/C expression increases with age and disease. In normal aging, the accumulation of prelamin-A/C is caused in part by the down-regulation of ZMPSTE24/FACE1 in response to oxidative stress.

Lamin A/C Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Lamin A/C Antibody - Images



Western blot analysis of Lamin A/C expression in HeLa cell treated with staurosporine.