

NEK7 Antibody
Rabbit mAb
Catalog # AP92261

Specification

NEK7 Antibody - Product Information

Application	WB, FC
Primary Accession	Q8TDX7
Reactivity	Rat
Clonality	Monoclonal
Other Names	
NEK7; NimA related protein kinase 7;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	34551 Da

NEK7 Antibody - Additional Information

Dilution	WB~~1:1000 FC~~1:10~50
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human NEK7
Description	Protein kinase which plays an important role in mitotic cell cycle progression. Required for microtubule nucleation activity of the centrosome, robust mitotic spindle formation and cytokinesis. Phosphorylates RPS6KB1.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

NEK7 Antibody - Protein Information

Name NEK7 {ECO:0000303|PubMed:11701951, ECO:0000312|HGNC:HGNC:13386}

Function

Protein kinase which plays an important role in mitotic cell cycle progression (PubMed:17101132, PubMed:19941817, PubMed:31409757). Required for microtubule nucleation activity of the centrosome, robust mitotic spindle formation and cytokinesis (PubMed:17586473, PubMed:19414596, PubMed:19941817).

target="_blank">19941817, PubMed:26522158, PubMed:31409757). Phosphorylates EML4 at 'Ser-146', promoting its dissociation from microtubules during mitosis which is required for efficient chromosome congression (PubMed:31409757). Phosphorylates RPS6KB1 (By similarity). Acts as an essential activator of the NLRP3 inflammasome assembly independently of its kinase activity (PubMed:26642356, PubMed:36442502, PubMed:39173637). Acts by unlocking NLRP3 following NLRP3 translocation into the microtubule organizing center (MTOC), relieving NLRP3 autoinhibition and promoting formation of the NLRP3:PYCARD complex, and activation of CASP1 (PubMed:26642356, PubMed:31189953, PubMed:36442502, PubMed:39173637). Serves as a cellular switch that enforces mutual exclusivity of the inflammasome response and cell division: interaction with NEK9 prevents interaction with NLRP3 and activation of the inflammasome during mitosis (PubMed:26642356, PubMed:31189953).

Cellular Location

Nucleus {ECO:0000250|UniProtKB:Q9ES74}. Cytoplasm. Cytoplasm, cytoskeleton, spindle pole. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Note=Present at centrosome throughout the cell cycle (PubMed:17586473). Also detected at spindle midzone of the anaphase cells and eventually concentrates at the midbody (PubMed:17586473). Interaction with ANKS3 prevents its translocation to the nucleus (By similarity). {ECO:0000250|UniProtKB:Q9ES74, ECO:0000269|PubMed:17586473}

Tissue Location


Highly expressed in lung, muscle, testis, brain, heart, liver, leukocyte and spleen. Lower expression in ovary, prostate and kidney. No expression seen in small intestine


NEK7 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NEK7 Antibody - Images





Western blot analysis of NEK7 expression in Jurkat cell lysate.