

**Hamartin Antibody**  
**Rabbit mAb**  
**Catalog # AP92337****Specification****Hamartin Antibody - Product Information**

Application	WB, IHC, FC, ICC
Primary Accession	<a href="#">Q92574</a>
Clonality	Monoclonal
<b>Other Names</b>	
Hamartin; kiaa0243; LAM; TSC; Tsc1; Tuberous sclerosis 1;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	129767 Da

**Hamartin Antibody - Additional Information**

Dilution	WB~~1:1000 IHC~~1:100~500 FC~~1:10~50 ICC~~N/A
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Hamartin
Description	In complex with TSC2, inhibits the nutrient-mediated or growth factor-stimulated phosphorylation of S6K1 and EIF4EBP1 by negatively regulating mTORC1 signaling. Seems not to be required for TSC2 GAP activity towards RHEB. Implicated as a tumor suppressor. Involved in microtubule-mediated protein transport, but this seems to be due to unregulated mTOR signaling.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

**Hamartin Antibody - Protein Information****Name** TSC1 {ECO:0000303|PubMed:9242607, ECO:0000312|HGNC:HGNC:12362}**Function**

Non-catalytic component of the TSC-TBC complex, a multiprotein complex that acts as a negative regulator of the canonical mTORC1 complex, an evolutionarily conserved central nutrient sensor that stimulates anabolic reactions and macromolecule biosynthesis to promote cellular biomass

generation and growth (PubMed:<a href="http://www.uniprot.org/citations/12172553" target="\_blank">12172553</a>, PubMed:<a href="http://www.uniprot.org/citations/12271141" target="\_blank">12271141</a>, PubMed:<a href="http://www.uniprot.org/citations/12906785" target="\_blank">12906785</a>, PubMed:<a href="http://www.uniprot.org/citations/15340059" target="\_blank">15340059</a>, PubMed:<a href="http://www.uniprot.org/citations/24529379" target="\_blank">24529379</a>, PubMed:<a href="http://www.uniprot.org/citations/28215400" target="\_blank">28215400</a>). The TSC-TBC complex acts as a GTPase-activating protein (GAP) for the small GTPase RHEB, a direct activator of the protein kinase activity of mTORC1 (PubMed:<a href="http://www.uniprot.org/citations/12906785" target="\_blank">12906785</a>, PubMed:<a href="http://www.uniprot.org/citations/15340059" target="\_blank">15340059</a>, PubMed:<a href="http://www.uniprot.org/citations/24529379" target="\_blank">24529379</a>). In absence of nutrients, the TSC-TBC complex inhibits mTORC1, thereby preventing phosphorylation of ribosomal protein S6 kinase (RPS6KB1 and RPS6KB2) and EIF4EBP1 (4E-BP1) by the mTORC1 signaling (PubMed:<a href="http://www.uniprot.org/citations/12271141" target="\_blank">12271141</a>, PubMed:<a href="http://www.uniprot.org/citations/24529379" target="\_blank">24529379</a>, PubMed:<a href="http://www.uniprot.org/citations/28215400" target="\_blank">28215400</a>, PubMed:<a href="http://www.uniprot.org/citations/33215753" target="\_blank">33215753</a>). The TSC-TBC complex is inactivated in response to nutrients, relieving inhibition of mTORC1 (PubMed:<a href="http://www.uniprot.org/citations/12172553" target="\_blank">12172553</a>, PubMed:<a href="http://www.uniprot.org/citations/24529379" target="\_blank">24529379</a>). Within the TSC-TBC complex, TSC1 stabilizes TSC2 and prevents TSC2 self-aggregation (PubMed:<a href="http://www.uniprot.org/citations/10585443" target="\_blank">10585443</a>, PubMed:<a href="http://www.uniprot.org/citations/28215400" target="\_blank">28215400</a>). Acts as a tumor suppressor (PubMed:<a href="http://www.uniprot.org/citations/9242607" target="\_blank">9242607</a>). Involved in microtubule-mediated protein transport via its ability to regulate mTORC1 signaling (By similarity). Also acts as a co-chaperone for HSP90AA1 facilitating HSP90AA1 chaperoning of protein clients such as kinases, TSC2 and glucocorticoid receptor NR3C1 (PubMed:<a href="http://www.uniprot.org/citations/29127155" target="\_blank">29127155</a>). Increases ATP binding to HSP90AA1 and inhibits HSP90AA1 ATPase activity (PubMed:<a href="http://www.uniprot.org/citations/29127155" target="\_blank">29127155</a>). Competes with the activating co-chaperone AHSA1 for binding to HSP90AA1, thereby providing a reciprocal regulatory mechanism for chaperoning of client proteins (PubMed:<a href="http://www.uniprot.org/citations/29127155" target="\_blank">29127155</a>). Recruits TSC2 to HSP90AA1 and stabilizes TSC2 by preventing the interaction between TSC2 and ubiquitin ligase HERC1 (PubMed:<a href="http://www.uniprot.org/citations/16464865" target="\_blank">16464865</a>, PubMed:<a href="http://www.uniprot.org/citations/29127155" target="\_blank">29127155</a>).

### Cellular Location

Lysosome membrane; Peripheral membrane protein. Cytoplasm, cytosol Note=Recruited to lysosomal membranes in a RHEB-dependent process in absence of nutrients (PubMed:24529379). In response to nutrients, the complex dissociates from lysosomal membranes and relocates to the cytosol (PubMed:24529379).

### Tissue Location


Highly expressed in skeletal muscle, followed by heart, brain, placenta, pancreas, lung, liver and kidney (PubMed:9242607). Also expressed in embryonic kidney cells (PubMed:9242607).

### Hamartin Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Hamartin Antibody - Images**

Western blot analysis of Hamartin expression in HeLa cell lysate.