

BACH1/BRIP1 Antibody
Rabbit mAb
Catalog # AP92369**Specification****BACH1/BRIP1 Antibody - Product Information**

Application	WB
Primary Accession	Q9BX63
Clonality	Monoclonal
Other Names	
BACH 1; BRIP 1; BRIP1; FANCI;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	140867 Da

BACH1/BRIP1 Antibody - Additional Information

Dilution	WB~~1:1000
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human BACH1/BRIP1
Description	DNA-dependent ATPase and 5' to 3' DNA helicase required for the maintenance of chromosomal stability. Acts late in the Fanconi anemia pathway, after FANCD2 ubiquitination. Involved in the repair of DNA double-strand breaks by homologous recombination in a manner that depends on its association with BRCA1.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

BACH1/BRIP1 Antibody - Protein Information**Name** BRIP1 ([HGNC:20473](#))**Function**

DNA-dependent ATPase and 5'-3' DNA helicase required for the maintenance of chromosomal stability (PubMed: [11301010](http://www.uniprot.org/citations/11301010) target="_blank">11301010, PubMed: [14983014](http://www.uniprot.org/citations/14983014) target="_blank">14983014, PubMed: [16116421](http://www.uniprot.org/citations/16116421) target="_blank">16116421, PubMed: [16153896](http://www.uniprot.org/citations/16153896) target="_blank">16153896, PubMed: [17596542](http://www.uniprot.org/citations/17596542) target="_blank">17596542, PubMed: [36608669](http://www.uniprot.org/citations/36608669) target="_blank">36608669). Acts late in the Fanconi anemia pathway, after FANCD2

ubiquitination (PubMed:14983014, PubMed:16153896). Involved in the repair of DNA double-strand breaks by homologous recombination in a manner that depends on its association with BRCA1 (PubMed:14983014, PubMed:16153896). Involved in the repair of abasic sites at replication forks by promoting the degradation of DNA-protein cross-links: acts by catalyzing unfolding of HMCES DNA-protein cross-link via its helicase activity, exposing the underlying DNA and enabling cleavage of the DNA- protein adduct by the SPRTN metalloprotease (PubMed:16116421, PubMed:36608669). Can unwind RNA:DNA substrates (PubMed:14983014). Unwinds G-quadruplex DNA; unwinding requires a 5'-single stranded tail (PubMed:18426915, PubMed:20639400).

Cellular Location

Nucleus. Cytoplasm

Tissue Location


Ubiquitously expressed, with highest levels in testis.

BACH1/BRIP1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

BACH1/BRIP1 Antibody - Images



Western blot analysis of BACH1/BRIP1 expression in HeLa cell lysate.