

Pumilio 2 Antibody
Rabbit mAb
Catalog # AP92379**Specification**

Pumilio 2 Antibody - Product Information

Application	WB, FC, ICC, IP
Primary Accession	Q8TB72
Reactivity	Rat
Clonality	Monoclonal
Other Names	
PUM2; PUMH2; Pumilio2; PUML2;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	114216 Da

Pumilio 2 Antibody - Additional Information

Dilution	WB~~1:1000 FC~~1:10~50 ICC~~N/A IP~~N/A
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Pumilio 2
Description	Sequence-specific RNA-binding protein that regulates translation and mRNA stability by binding the 3'-UTR of mRNA targets. Its interactions and tissue specificity suggest that it may be required to support proliferation and self-renewal of stem cells by regulating the translation of key transcripts.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Pumilio 2 Antibody - Protein Information**Name** PUM2**Synonyms** KIAA0235, PUMH2**Function**

Sequence-specific RNA-binding protein that acts as a post- transcriptional repressor by binding the 3'-UTR of mRNA targets. Binds to an RNA consensus sequence, the Pumilio Response Element

(PRE), 5'- UGUANAUA-3', that is related to the Nanos Response Element (NRE) ([PubMed:21397187](http://www.uniprot.org/citations/21397187)). Mediates post-transcriptional repression of transcripts via different mechanisms: acts via direct recruitment of the CCR4-POP2-NOT deadenylase leading to translational inhibition and mRNA degradation ([PubMed:22955276](http://www.uniprot.org/citations/22955276)). Also mediates deadenylation- independent repression by promoting accessibility of miRNAs ([PubMed:18776931](http://www.uniprot.org/citations/18776931), [PubMed:22345517](http://www.uniprot.org/citations/22345517)). Acts as a post-transcriptional repressor of E2F3 mRNAs by binding to its 3'-UTR and facilitating miRNA regulation ([PubMed:22345517](http://www.uniprot.org/citations/22345517)). Plays a role in cytoplasmic sensing of viral infection ([PubMed:25340845](http://www.uniprot.org/citations/25340845)). Represses a program of genes necessary to maintain genomic stability such as key mitotic, DNA repair and DNA replication factors. Its ability to repress those target mRNAs is regulated by the lncRNA NORAD (non-coding RNA activated by DNA damage) which, due to its high abundance and multitude of PUMILIO binding sites, is able to sequester a significant fraction of PUM1 and PUM2 in the cytoplasm ([PubMed:26724866](http://www.uniprot.org/citations/26724866)). May regulate DCUN1D3 mRNA levels ([PubMed:25349211](http://www.uniprot.org/citations/25349211)). May support proliferation and self-renewal of stem cells. Binds specifically to miRNA MIR199A precursor, with PUM1, regulates miRNA MIR199A expression at a postranscriptional level ([PubMed:28431233](http://www.uniprot.org/citations/28431233)).

Cellular Location

Cytoplasm. Cytoplasmic granule. Cytoplasm, perinuclear region. Note=The cytoplasmic granules are stress granules which are a dense aggregation in the cytosol composed of proteins and RNAs that appear when the cell is under stress. Colocalizes with NANOS3 in the stress granules Colocalizes with NANOS1 and SNAPIN in the perinuclear region of germ cells.

Tissue Location


Expressed in male germ cells of adult testis (at protein level). Highly expressed in testis and ovary. Predominantly expressed in stem cells and germ cells. Expressed at lower level in brain, heart, kidney, liver, muscle, placenta, intestine and stomach Expressed in cerebellum, corpus callosum, caudate nucleus, hippocampus, medulla oblongata and putamen. Expressed in all fetal tissues tested


Pumilio 2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Pumilio 2 Antibody - Images





Western blot analysis of Pumilio 2 expression in 293T cell lysate.