

**ADA Antibody  
Rabbit mAb  
Catalog # AP92392**

## Specification

**ADA Antibody - Product Information**

Application	WB, ICC, IP
Primary Accession	<a href="#">P00813</a>
Reactivity	Rat
Clonality	Monoclonal
<b>Other Names</b>	
ada; ADA1;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	40764 Da

#### **ADA Antibody - Additional Information**

Dilution	WB~~1:1000 ICC~~N/A IP~~N/A
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human ADA
Description	Catalyzes the hydrolytic deamination of adenosine and 2-deoxyadenosine. Plays an important role in purine metabolism and in adenosine homeostasis. Modulates signaling by extracellular adenosine, and so contributes indirectly to cellular signaling events.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

## **ADA Antibody - Protein Information**

Name ADA

## Synonyms ADA1

## Function

[8452534](http://www.uniprot.org/citations/8452534), PubMed:[9361033](http://www.uniprot.org/citations/9361033)). Plays an important role in purine metabolism and in adenosine homeostasis. Modulates signaling by extracellular adenosine, and so contributes indirectly to cellular signaling events. Acts as a positive regulator of T-cell coactivation, by binding DPP4 (PubMed:[20959412](http://www.uniprot.org/citations/20959412)). Its interaction with DPP4 regulates lymphocyte-epithelial cell adhesion (PubMed:[11772392](http://www.uniprot.org/citations/11772392)). Enhances dendritic cell immunogenicity by affecting dendritic cell costimulatory molecule expression and cytokines and chemokines secretion (By similarity). Enhances CD4+ T-cell differentiation and proliferation (PubMed:[20959412](http://www.uniprot.org/citations/20959412)). Acts as a positive modulator of adenosine receptors ADORA1 and ADORA2A, by enhancing their ligand affinity via conformational change (PubMed:[23193172](http://www.uniprot.org/citations/23193172)). Stimulates plasminogen activation (PubMed:[15016824](http://www.uniprot.org/citations/15016824)). Plays a role in male fertility (PubMed:[21919946](http://www.uniprot.org/citations/21919946), PubMed:[26166670](http://www.uniprot.org/citations/26166670)). Plays a protective role in early postimplantation embryonic development (By similarity). Also responsible for the deamination of cordycepin (3'-deoxyadenosine), a fungal natural product that shows antitumor, antibacterial, antifungal, antivirus, and immune regulation properties (PubMed:[26038697](http://www.uniprot.org/citations/26038697)).

## Cellular Location

Cell membrane; Peripheral membrane protein; Extracellular side. Cell junction. Cytoplasmic vesicle lumen {ECO:0000250|UniProtKB:P03958}. Cytoplasm. Lysosome. Note=Colocalized with DPP4 at the cell surface.

## Tissue Location


Found in all tissues, occurs in large amounts in T- lymphocytes (PubMed:20959412). Expressed at the time of weaning in gastrointestinal tissues.

## **ADA Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
  - Blocking Peptides
  - Dot Blot
  - Immunohistochemistry
  - Immunofluorescence
  - Immunoprecipitation
  - Flow Cytometry
  - Cell Culture

## **ADA Antibody - Images**



Western blot analysis of ADA expression in Jurkat cell lysate.