

### **ALKBH1 Antibody**

Rabbit mAb Catalog # AP92535

#### **Specification**

### **ALKBH1 Antibody - Product Information**

Application WB, IHC, IP
Primary Accession Q13686
Clonality Monoclonal

**Other Names** 

ABH; ABH1; alkB; ALKBH; ALKBH1; hABH;

Isotype Rabbit IgG
Host Rabbit
Calculated MW 43832 Da

## **ALKBH1 Antibody - Additional Information**

Dilution WB~~1:1000

IHC~~1:100~500

IP~~N/A

Purification Affinity-chromatography

Immunogen A synthesized peptide derived from human

ALKRH1

Description Dioxygenase that repairs alkylated

single-stranded DNA and RNA containing

3-methylcytosine by oxidative

demethylation. Requires molecular oxygen,

alpha-ketoglutarate and iron.

Storage Condition and Buffer Rabbit IgG in phosphate buffered saline ,

pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid

freeze / thaw cycle.

### **ALKBH1 Antibody - Protein Information**

Name ALKBH1 (HGNC:17911)

#### **Function**

Dioxygenase that acts on nucleic acids, such as DNA and tRNA (PubMed:<a

href="http://www.uniprot.org/citations/18603530" target="\_blank">18603530</a>, PubMed:<a href="http://www.uniprot.org/citations/27497299" target="\_blank">27497299</a>, PubMed:<a href="http://www.uniprot.org/citations/27745969" target="\_blank">27745969</a>). Requires molecular oxygen, alpha-ketoglutarate and iron (PubMed:<a

href="http://www.uniprot.org/citations/18603530" target="\_blank">18603530</a>, PubMed:<a href="http://www.uniprot.org/citations/27497299" target="\_blank">27497299</a>). A number of activities have been described for this dioxygenase, but recent results suggest that it mainly acts on tRNAs and mediates their demethylation or oxidation depending on the context and subcellular



Tel: 858.875.1900 Fax: 858.875.1999

compartment (PubMed:<a href="http://www.uniprot.org/citations/27497299" target=" blank">27497299</a>, PubMed:<a href="http://www.uniprot.org/citations/27745969" target="blank">27745969</a>). Mainly acts as a tRNA demethylase by removing N(1)-methyladenine from various tRNAs, with a preference for N(1)-methyladenine at position 58 (m1A58) present on a stem loop structure of tRNAs (PubMed:<a href="http://www.uniprot.org/citations/27745969" target=" blank">27745969</a>). Acts as a regulator of translation initiation and elongation in response to glucose deprivation: regulates both translation initiation, by mediating demethylation of tRNA(Met), and translation elongation, N(1)methyladenine-containing tRNAs being preferentially recruited to polysomes to promote translation elongation (PubMed: <a href="http://www.uniprot.org/citations/27745969" target=" blank">27745969</a>). In mitochondrion, specifically interacts with mt-tRNA(Met) and mediates oxidation of mt-tRNA(Met) methylated at cytosine(34) to form 5- formylcytosine (f(5)c) at this position (PubMed: <a href="http://www.uniprot.org/citations/27497299" target=" blank">27497299</a>), mt-tRNA(Met) containing the f(5)c modification at the wobble position enables recognition of the AUA codon in addition to the AUG codon, expanding codon recognition in mitochondrial translation (PubMed:<a href="http://www.uniprot.org/citations/27497299" target=" blank">27497299</a>). Specifically demethylates DNA methylated on the 6th position of adenine (N(6)-methyladenosine) DNA (PubMed:<a href="http://www.uniprot.org/citations/30017583" target=" blank">30017583</a>, PubMed:<a href="http://www.uniprot.org/citations/30392959" target=" blank">30392959</a>). N(6)- methyladenosine (m6A) DNA is present at some L1 elements in embryonic stem cells and probably promotes their silencing (By similarity). Demethylates mRNAs containing N(3)-methylcytidine modification (PubMed: <a href="http://www.uniprot.org/citations/31188562" target=" blank">31188562</a>). Also able to repair alkylated single-stranded DNA by oxidative demethylation, but with low activity (PubMed:<a href="http://www.uniprot.org/citations/18603530" target=" blank">18603530</a>). Also has DNA lyase activity and introduces double-stranded breaks at abasic sites: cleaves both single-stranded DNA and double-stranded DNA at abasic sites, with the greatest activity towards double-stranded DNA with two abasic sites (PubMed: <a href="http://www.uniprot.org/citations/19959401" target=" blank">19959401</a>). DNA lyase activity does not require alpha-ketoglutarate and iron and leads to the formation of an irreversible covalent protein-DNA adduct with the 5' DNA product (PubMed: <a href="http://www.uniprot.org/citations/19959401" target=" blank">19959401</a>, PubMed:<a href="http://www.uniprot.org/citations/23577621" target="blank">23577621</a>). DNA lyase activity is not required during base excision repair and class switch recombination of the immunoglobulin heavy chain during B lymphocyte activation. May play a role in placental trophoblast lineage differentiation (By similarity).

## **Cellular Location**


Nucleus. Mitochondrion. Note=Mainly localizes in euchromatin, largely excluded from heterochromatin and nucleoli (By similarity). {ECO:0000250|UniProtKB:P0CB42}

**Tissue Location** Ubiquitous.

## **ALKBH1 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety





# • Cell Culture **ALKBH1 Antibody - Images**

