

WDR4 Antibody
Rabbit mAb
Catalog # AP92591**Specification****WDR4 Antibody - Product Information**

Application	WB, IHC, FC, ICC
Primary Accession	P57081
Clonality	Monoclonal
Other Names	
TRM82; WD repeat protein 4; wdr4;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	45490 Da

WDR4 Antibody - Additional Information

Dilution	WB~~1:1000 IHC~~1:100~500 FC~~1:10~50 ICC~~N/A
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human WDR4
Description	Required for the formation of N(7)-methylguanine at position 46 (m7G46) in tRNA. In the complex, it is required to stabilize and induce conformational change of the catalytic subunit.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

WDR4 Antibody - Protein Information**Name** WDR4**Function**

Non-catalytic component of the METTL1-WDR4 methyltransferase complex required for the formation of N(7)-methylguanine in a subset of RNA species, such as tRNAs, mRNAs and microRNAs (miRNAs) (PubMed:12403464, PubMed:31031083, PubMed:31031084, PubMed:36599982, PubMed:36599985)

target="_blank">36599985, PubMed:37369656). In the METTL1-WDR4 methyltransferase complex, WDR4 acts as a scaffold for tRNA-binding (PubMed:36599982, PubMed:36599985, PubMed:37369656). Required for the formation of N(7)-methylguanine at position 46 (m7G46) in a large subset of tRNAs that contain the 5'-RAGGU-3' motif within the variable loop (PubMed:12403464, PubMed:34352206, PubMed:34352207, PubMed:36599982, PubMed:36599985, PubMed:37369656). M7G46 interacts with C13-G22 in the D-loop to stabilize tRNA tertiary structure and protect tRNAs from decay (PubMed:36599982, PubMed:36599985). Also required for the formation of N(7)-methylguanine at internal sites in a subset of mRNAs (PubMed:31031084, PubMed:37379838). Also required for methylation of a specific subset of miRNAs, such as let-7 (PubMed:31031083). Independently of METTL1, also plays a role in genome stability: localizes at the DNA replication site and regulates endonucleolytic activities of FEN1 (PubMed:26751069).

Cellular Location


Nucleus. Chromosome Note=Localizes at the site of nascent DNA synthesis


WDR4 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

WDR4 Antibody - Images





Western blot analysis of WDR4 expression in HepG2 cell lysate.