

## **Complement factor B Antibody**

Rabbit mAb Catalog # AP92620

### **Specification**

## **Complement factor B Antibody - Product Information**

Application WB
Primary Accession P00751
Clonality Monoclonal

**Other Names** 

C3/C5 convertase; Complement factor B; PBF2; Properdin factor B;

Isotype Rabbit IgG
Host Rabbit
Calculated MW 85533 Da

# **Complement factor B Antibody - Additional Information**

Dilution WB~~1:1000

Purification Affinity-chromatography

Immunogen A synthesized peptide derived from human

**Complement factor B** 

Description Factor B which is part of the alternate

pathway of the complement system is cleaved by factor D into 2 fragments: Ba and Bb. Bb, a serine protease, then combines with complement factor 3b to

generate the C3 or C5 convertase.

Storage Condition and Buffer Rabbit IgG in phosphate buffered saline ,

pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid

freeze / thaw cycle.

# **Complement factor B Antibody - Protein Information**

Name CFB (HGNC:1037)

Synonyms BF, BFD

#### **Function**

Precursor of the catalytic component of the C3 and C5 convertase complexes of the alternative pathway of the complement system, a cascade of proteins that leads to phagocytosis and breakdown of pathogens and signaling that strengthens the adaptive immune system (PubMed:<a href="http://www.uniprot.org/citations/3638964" target="\_blank">3638964</a>, PubMed:<a href="http://www.uniprot.org/citations/624565" target="\_blank">624565</a>, PubMed:<a href="http://www.uniprot.org/citations/6554279" target="\_blank">6554279</a>, PubMed:<a href="http://www.uniprot.org/citations/6919543" target="\_blank">6919543</a>, PubMed:<a href="http://www.uniprot.org/citations/9748277" target="\_blank">9748277</a>). The alternative



complement pathway acts as an amplification loop that enhances other complement pathways (classical, lectin and GZMK) by promoting formation of additional C3 and C5 convertases (PubMed:<a href="http://www.uniprot.org/citations/3638964" target="\_blank">3638964</a>, PubMed:<a href="http://www.uniprot.org/citations/624565" target="\_blank">624565</a>, PubMed:<a href="http://www.uniprot.org/citations/6554279" target="\_blank">6554279</a>, PubMed:<a href="http://www.uniprot.org/citations/6919543" target="\_blank">6919543</a>, PubMed:<a href="http://www.uniprot.org/citations/9748277" target="\_blank">9748277</a>). CFB is cleaved and activated by CFD to generate Ba and Bb chains; Bb chain constituting the catalytic component of the C3 and C5 convertases (PubMed:<a href="http://www.uniprot.org/citations/6769474" target="\_blank">6769474</a>, PubMed:<a href="http://www.uniprot.org/citations/9748277" target="\_blank">9748277</a>).


**Cellular Location** Secreted.

## **Complement factor B Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

## Complement factor B Antibody - Images



Western blot analysis of Complement factor B expression in Human plasma lysate.