

UCHL3 Antibody

Rabbit mAb Catalog # AP92669

Specification

UCHL3 Antibody - Product Information

Application	WB, IHC
Primary Accession	<u>P15374</u>
Reactivity	Rat
Clonality	Monoclonal
Other Names	
Ubiquitin thioesterase L3; UCHL3;	
Isativaa	Rabbit IgG
lsotype Host	Rabbit
Calculated MW	26183 Da
	20105 Da

UCHL3 Antibody - Additional Information

Dilution	WB~~1:1000 IHC~~1:100~500
Purification Immunogen Description	Affinity-chromatography A synthesized peptide derived from UCHL3 Deubiquitinating enzyme (DUB) that controls levels of cellular ubiquitin through processing of ubiquitin precursors and ubiquitinated proteins. Thiol protease that recognizes and hydrolyzes a peptide bond at the C-terminal glycine of either ubiquitin or NEDD8.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

UCHL3 Antibody - Protein Information

Name UCHL3

Function

Deubiquitinating enzyme (DUB) that controls levels of cellular ubiquitin through processing of ubiquitin precursors and ubiquitinated proteins. Thiol protease that recognizes and hydrolyzes a peptide bond at the C-terminal glycine of either ubiquitin or NEDD8. Has a 10-fold preference for Arg and Lys at position P3", and exhibits a preference towards 'Lys-48'-linked ubiquitin chains. Deubiquitinates ENAC in apical compartments, thereby regulating apical membrane recycling. Indirectly increases the phosphorylation of IGFIR, AKT and FOXO1 and promotes insulin-signaling and insulin-induced adipogenesis. Required for stress-response retinal, skeletal muscle and germ cell maintenance. May be involved in working memory. Can hydrolyze UBB(+1), a mutated form of



ubiquitin which is not effectively degraded by the proteasome and is associated with neurogenerative disorders.

Cellular Location Cytoplasm.

Tissue Location Highly expressed in heart, skeletal muscle, and testis.

UCHL3 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

UCHL3 Antibody - Images



Western blot analysis of UCHL3 expression in K562 cell lysate.