

## **ACTR5 Antibody (C-term)**

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP9266b

## **Specification**

# **ACTR5 Antibody (C-term) - Product Information**

Application IHC-P, WB,E Primary Accession Q9H9F9

Reactivity Human, Hamster

Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Calculated MW 68297
Antigen Region 539-568

## **ACTR5 Antibody (C-term) - Additional Information**

#### **Gene ID** 79913

### **Other Names**

Actin-related protein 5, hARP5, Sarcoma antigen NY-SAR-16, ACTR5, ARP5

# Target/Specificity

This ACTR5 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 539-568 amino acids from the C-terminal region of human ACTR5.

#### **Dilution**

IHC-P~~1:50~100 WB~~1:1000

E~~Use at an assay dependent concentration.

### **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

## **Precautions**

ACTR5 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## **ACTR5 Antibody (C-term) - Protein Information**

### Name ACTR5

**Synonyms** ARP5





**Function** Proposed core component of the chromatin remodeling INO80 complex which is involved in transcriptional regulation, DNA replication and probably DNA repair. Involved in DNA double-strand break repair and UV-damage excision repair.

#### **Cellular Location**


Nucleus. Cytoplasm. Note=Predominantly nuclear but undergoes nucleo-cytoplasmic shuttling (PubMed:19014934). Localized to interphase nuclei, but not nucleoli; excluded from chromosomes as mitosis progresses (PubMed:18163988)

# **ACTR5 Antibody (C-term) - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

## **ACTR5 Antibody (C-term) - Images**



Western blot analysis of ACTR5 Antibody (C-term) (Cat. #AP9266b) in CHO cell line lysates (35ug/lane). ACTR5 (arrow) was detected using the purified Pab.







Formalin-fixed and paraffin-embedded human brain tissue reacted with ACTR5 Antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

# ACTR5 Antibody (C-term) - Background

ACTR5 may be involved in transcription regulation via its interaction with the INO80 complex, a chromatin remodeling complex

## **ACTR5 Antibody (C-term) - References**

Kitayama, K., et.al., Exp. Cell Res. 315 (2), 206-217 (2009) Jin, J., et.al., J. Biol. Chem. 280 (50), 41207-41212 (2005) Lee, S.Y., et.al., Proc. Natl. Acad. Sci. U.S.A. 100 (5), 2651-2656 (2003)