

GBP1 Antibody

Rabbit mAb Catalog # AP92681

Specification

GBP1 Antibody - Product Information

Application WB, IHC, ICC Primary Accession P32455
Clonality Monoclonal

Other Names

GBP1; GTP binding protein 1; Guanine nucleotide binding protein 1; Guanylate binding protein 1; HuGBP1; Interferon induced guanylate binding protein 1;

Isotype Rabbit IgG
Host Rabbit
Calculated MW 67931 Da

GBP1 Antibody - Additional Information

Dilution WB~~1:1000

IHC~~1:100~500

ICC~~N/A

Purification Affinity-chromatography

Immunogen A synthesized peptide derived from human

GBP1

Description Binds GTP, GDP and GMP.

Storage Condition and Buffer Rabbit IgG in phosphate buffered saline ,

pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid

freeze / thaw cycle.

GBP1 Antibody - Protein Information

Name GBP1 {ECO:0000303|PubMed:7512561, ECO:0000312|HGNC:HGNC:4182}

Function

Interferon (IFN)-inducible GTPase that plays important roles in innate immunity against a diverse range of bacterial, viral and protozoan pathogens (PubMed:16511497, PubMed:22106366, PubMed:2914452, PubMed:31268602, PubMed:32510692, PubMed:32581219, PubMed:37797010, PubMed:7512561, PubMed:7512561, PubMed:7512561, PubMed:<a href="http://www.uniprot.org/citations/16511497"



target="_blank">16511497, PubMed:32510692, PubMed:7512561). Following infection, recruited to the pathogen-containing vacuoles or vacuole-escaped bacteria and promotes both inflammasome assembly and autophagy (PubMed:29144452, PubMed:31268602). Acts as a positive regulator of inflammasome assembly by facilitating the detection of inflammasome ligands from pathogens (PubMed:31268602, PubMed:32510692, PubMed:32581219). Involved in the lysis of pathogen-containing vacuoles, releasing pathogens into the cytosol (By similarity). Following pathogen release in the cytosol, forms a protein coat in a GTPase-dependent manner that encapsulates pathogens and promotes the detection of ligands by pattern recognition receptors (PubMed:<a href="http://www.uniprot.org/citations/32510692"

target="_blank">32510692, PubMed:32581219). Plays a key role in inflammasome assembly in response to infection by Gram-negative bacteria: following pathogen release in the cytosol, forms a protein coat that encapsulates Gram-negative bacteria and directly binds to lipopolysaccharide (LPS), disrupting the O-antigen barrier and unmasking lipid A that is that detected by the non-canonical inflammasome effector CASP4/CASP11 (PubMed:32510692, PubMed:32581219). Also promotes recruitment of proteins that mediate bacterial cytolysis, leading to release double-stranded DNA (dsDNA) that activates the AIM2 inflammasome (PubMed:31268602). Involved in autophagy by regulating bacteriolytic peptide generation via its interaction with ubiquitin-binding protein SQSTM1, which delivers monoubiquitinated proteins to autolysosomes for the generation of bacteriolytic peptides (By similarity). Confers protection to several pathogens, including the bacterial pathogens L.monocytogenes and M.bovis BCG as well as the protozoan pathogen T.gondii (PubMed:<a href="http://www.uniprot.org/citations/31268602"

target="_blank">31268602). Exhibits antiviral activity against influenza virus (PubMed:22106366).

Cellular Location

Cytoplasmic vesicle membrane; Lipid-anchor; Cytoplasmic side. Golgi apparatus membrane; Lipid-anchor; Cytoplasmic side. Cytoplasmic side.

GBP1 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- Immunoprecipitation



- Flow Cytomety
- Cell Culture

GBP1 Antibody - Images



Western blot analysis of GBP1 expression in HeLa cell lysate.