

**USP39 Antibody**  
**Rabbit mAb**  
**Catalog # AP92705****Specification****USP39 Antibody - Product Information**

Application	WB, IHC, ICC
Primary Accession	<a href="#">Q53GS9</a>
Reactivity	Rat
Clonality	Monoclonal
<b>Other Names</b>	
Inactive ubiquitin specific peptidase 39; SAD1; snRNP ASSEMBLY DEFECTIVE 1; SNRNP65; Ubiquitin specific peptidase 39; USP39;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	65381 Da

**USP39 Antibody - Additional Information**

Dilution	WB~~1:1000 IHC~~1:100~500 ICC~~N/A
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human USP39
Description	May play a role in mRNA splicing. It is unsure if the protein really exhibits hydrolase activity. Could be a competitor of ubiquitin C-terminal hydrolases (UCHs).
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

**USP39 Antibody - Protein Information****Name** USP39 ([HGNC:20071](#))**Function**

Deubiquitinating enzyme that plays a role in many cellular processes including cellular antiviral response, epithelial morphogenesis, DNA repair or B-cell development (PubMed:<a href="http://www.uniprot.org/citations/33127822" target="\_blank">33127822</a>, PubMed:<a href="http://www.uniprot.org/citations/34614178" target="\_blank">34614178</a>). Plays a role in pre-mRNA splicing as a component of the U4/U6-U5 tri-snRNP, one of the building blocks of the precatalytic spliceosome (PubMed:<a href="http://www.uniprot.org/citations/11350945" target="\_blank">11350945</a>, PubMed:<a href="http://www.uniprot.org/citations/26912367" target="\_blank">26912367</a>). Specifically regulates immunoglobulin gene rearrangement in a

spliceosome-dependent manner, which involves modulating chromatin interactions at the Igh locus and therefore plays an essential role in B-cell development (By similarity). Regulates AURKB mRNA levels, and thereby plays a role in cytokinesis and in the spindle checkpoint (PubMed:<a href="http://www.uniprot.org/citations/18728397" target="\_blank">18728397</a>). Regulates apoptosis and G2/M cell cycle checkpoint in response to DNA damage by deubiquitinating and stabilizing CHK2 (PubMed:<a href="http://www.uniprot.org/citations/30771428" target="\_blank">30771428</a>). Also plays an important role in DNA repair by controlling the recruitment of XRCC4/LIG4 to DNA double-strand breaks for non-homologous end-joining repair (PubMed:<a href="http://www.uniprot.org/citations/34614178" target="\_blank">34614178</a>). Participates in antiviral activity by affecting the type I IFN signaling by stabilizing STAT1 and decreasing its 'Lys-6'-linked ubiquitination (PubMed:<a href="http://www.uniprot.org/citations/33127822" target="\_blank">33127822</a>). Contributes to non-canonical Wnt signaling during epidermal differentiation (By similarity). Acts as a negative regulator NF-kappa-B activation through deubiquitination of 'Lys-48'-linked ubiquitination of NFKBIA (PubMed:<a href="http://www.uniprot.org/citations/36651806" target="\_blank">36651806</a>).

### Cellular Location


Nucleus

### USP39 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### USP39 Antibody - Images



Western blot analysis of USP39 expression in HeLa cell lysate.