

**Osteoprotegerin Antibody**  
**Rabbit mAb**  
**Catalog # AP92717****Specification**

---

**Osteoprotegerin Antibody - Product Information**

Application	IHC, ICC
Primary Accession	<a href="#">O00300</a>
Clonality	Monoclonal
<b>Other Names</b>	
OCIF; OPG; Osteoclastogenesis inhibitory factor; Osteoprotegerin; PDB5; TNFRSF11B; TR1;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	46026 Da

**Osteoprotegerin Antibody - Additional Information**

Dilution	IHC~~1:100~500 ICC~~N/A
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Osteoprotegerin
Description	Acts as decoy receptor for RANKL and thereby neutralizes its function in osteoclastogenesis. Inhibits the activation of osteoclasts and promotes osteoclast apoptosis in vitro. Bone homeostasis seems to depend on the local RANKL/OPG ratio. May also play a role in preventing arterial calcification. May act as decoy receptor for TRAIL and protect against apoptosis. TRAIL binding blocks the inhibition of osteoclastogenesis.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

**Osteoprotegerin Antibody - Protein Information****Name** TNFRSF11B**Synonyms** OCIF, OPG**Function**

Acts as a decoy receptor for TNFSF11/RANKL and thereby neutralizes its function in osteoclastogenesis. Inhibits the activation of osteoclasts and promotes osteoclast apoptosis in

vitro. Bone homeostasis seems to depend on the local ratio between TNFSF11 and TNFRSF11B. May also play a role in preventing arterial calcification. May act as decoy receptor for TNFSF10/TRAIL and protect against apoptosis. TNFSF10/TRAIL binding blocks the inhibition of osteoclastogenesis.

**Cellular Location**

Secreted.

**Tissue Location**

Highly expressed in adult lung, heart, kidney, liver, spleen, thymus, prostate, ovary, small intestine, thyroid, lymph node, trachea, adrenal gland, testis, and bone marrow. Detected at very low levels in brain, placenta and skeletal muscle. Highly expressed in fetal kidney, liver and lung

**Osteoprotegerin Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Osteoprotegerin Antibody - Images**