

ATP1B2 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP9271c**Specification**

ATP1B2 Antibody (Center) - Product Information

Application	WB, FC, IHC-P,E
Primary Accession	P14415
Other Accession	P13638 , Q8WMG3 , P14231 , Q28030
Reactivity	Human
Predicted	Bovine, Mouse, Rabbit, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	33367
Antigen Region	115-141

ATP1B2 Antibody (Center) - Additional Information**Gene ID** 482**Other Names**

Sodium/potassium-transporting ATPase subunit beta-2, Adhesion molecule in glia, AMOG,
Sodium/potassium-dependent ATPase subunit beta-2, ATP1B2

Target/Specificity

This ATP1B2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 115-141 amino acids from the Central region of human ATP1B2.

Dilution

WB~~1:1000
FC~~1:10~50
IHC-P~~1:50~100
E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ATP1B2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

ATP1B2 Antibody (Center) - Protein Information

Name ATP1B2

Function This is the non-catalytic component of the active enzyme, which catalyzes the hydrolysis of ATP coupled with the exchange of Na(+) and K(+) ions across the plasma membrane. The exact function of the beta-2 subunit is not known.


Cellular Location

Cell membrane; Single-pass type II membrane protein


ATP1B2 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ATP1B2 Antibody (Center) - Images

Western blot analysis of lysates from K562 cell line and rat liver tissue lysate(from left to right), using ATP1B2 Antibody (Center)(Cat. #AP9271c). AP9271c was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 35ug per lane.



Formalin-fixed and paraffin-embedded human hepatocarcinoma reacted with ATP1B2 Antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



ATP1B2 Antibody (Center) (Cat. #AP9271c) flow cytometry analysis of K562 cells (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

ATP1B2 Antibody (Center) - Background

The protein belongs to the family of Na⁺/K⁺ and H⁺/K⁺ ATPases beta chain proteins, and to the subfamily of Na⁺/K⁺ -ATPases. Na⁺/K⁺ -ATPase is an integral membrane protein responsible for establishing and maintaining the electrochemical gradients of Na and K ions across the plasma membrane. These gradients are essential for osmoregulation, for sodium-coupled transport of a variety of organic and inorganic molecules, and for electrical excitability of nerve and muscle. This enzyme is composed of two subunits, a large catalytic subunit (alpha) and a smaller glycoprotein subunit (beta). The beta subunit regulates, through assembly of alpha/beta heterodimers, the number of sodium pumps transported to the plasma membrane.

ATP1B2 Antibody (Center) - References

- Guey, L.T., et.al., Eur. Urol. 57 (2), 283-292 (2010)
- Tokhtaeva, E., et.al., Biochemistry 48 (48), 11421-11431 (2009)
- Hosgood, H.D. et.al., Respir Med 103 (12), 1866-1870 (2009)