

ZNF259 Antibody
Rabbit mAb
Catalog # AP92733

Specification

ZNF259 Antibody - Product Information

Application	WB, IHC
Primary Accession	O75312
Reactivity	Rat
Clonality	Monoclonal
Other Names	
ZNF259; ZPR1;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	50925 Da

ZNF259 Antibody - Additional Information

Dilution	WB~~1:1000 IHC~~1:100~500
Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human ZNF259
Description	May be a signaling molecule that communicates mitogenic signals from the cytoplasm to the nucleus.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

ZNF259 Antibody - Protein Information

Name ZPR1

Synonyms ZNF259

Function

Acts as a signaling molecule that communicates proliferative growth signals from the cytoplasm to the nucleus. It is involved in the positive regulation of cell cycle progression (PubMed:29851065). Plays a role for the localization and accumulation of the survival motor neuron protein SMN1 in sub-nuclear bodies, including gems and Cajal bodies. Induces neuron differentiation and stimulates axonal growth and formation of growth cone in spinal cord motor neurons. Plays a role in the splicing of cellular pre-mRNAs. May be involved in H(2)O(2)-induced neuronal cell death.

Cellular Location

Nucleus. Nucleus, nucleolus. Nucleus, gem. Nucleus, Cajal body. Cytoplasm, perinuclear region. Cytoplasm. Cell projection, axon. Cell projection, growth cone. Note=Colocalized with SMN1 in Gemini of coiled bodies (gems), Cajal bodies, axon and growth cones of neurons (By similarity) Localized predominantly in the cytoplasm in serum-starved cells growth arrested in G0 of the mitotic cell cycle. Localized both in the nucleus and cytoplasm at the G1 phase of the mitotic cell cycle. Accumulates in the subnuclear bodies during progression into the S phase of the mitotic cell cycle. Diffusely localized throughout the cell during mitosis. Colocalized with NPAT and SMN1 in nuclear bodies including gems (Gemini of coiled bodies) and Cajal bodies in a cell cycle-dependent manner. Translocates together with EEF1A1 from the cytoplasm to the nucleolus after treatment with mitogens. Colocalized with EGFR in the cytoplasm of quiescent cells. Translocates from the cytoplasm to the nucleus in a epidermal growth factor (EGF)-dependent manner

Tissue Location


Expressed in fibroblast; weakly expressed in fibroblast of spinal muscular atrophy (SMA) patients

ZNF259 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZNF259 Antibody - Images



Western blot analysis of ZNF259 expression in MCF7 cell lysate.