

USP39 Antibody
Rabbit mAb
Catalog # AP92736**Specification****USP39 Antibody - Product Information**

Application **WB, IHC, ICC**
Primary Accession **[Q53GS9](#)**
Clonality **Monoclonal**
Other Names
CGI 21; SAD1; SNRNP65; USP39;

Isotype **Rabbit IgG**
Host **Rabbit**
Calculated MW **65381 Da**

USP39 Antibody - Additional Information

Dilution **WB~~1:1000**
IHC~~1:100~500
ICC~~N/A
Purification **Affinity-chromatography**
Immunogen **A synthesized peptide derived from human USP39**
Description **May play a role in mRNA splicing. It is unsure if the protein really exhibits hydrolase activity. Could be a competitor of ubiquitin C-terminal hydrolases (UCHs). Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.**
Storage Condition and Buffer

USP39 Antibody - Protein Information

Name USP39 ([HGNC:20071](#))

Function

Deubiquitinating enzyme that plays a role in many cellular processes including cellular antiviral response, epithelial morphogenesis, DNA repair or B-cell development (PubMed:33127822, PubMed:34614178). Plays a role in pre-mRNA splicing as a component of the U4/U6-U5 tri-snRNP, one of the building blocks of the precatalytic spliceosome (PubMed:11350945, PubMed:26912367). Specifically regulates immunoglobulin gene rearrangement in a spliceosome-dependent manner, which involves modulating chromatin interactions at the Igh locus and therefore plays an essential role in B-cell development (By similarity). Regulates AURKB

mRNA levels, and thereby plays a role in cytokinesis and in the spindle checkpoint (PubMed:18728397). Regulates apoptosis and G2/M cell cycle checkpoint in response to DNA damage by deubiquitinating and stabilizing CHK2 (PubMed:30771428). Also plays an important role in DNA repair by controlling the recruitment of XRCC4/LIG4 to DNA double-strand breaks for non-homologous end-joining repair (PubMed:34614178). Participates in antiviral activity by affecting the type I IFN signaling by stabilizing STAT1 and decreasing its 'Lys-6'-linked ubiquitination (PubMed:33127822). Contributes to non-canonical Wnt signaling during epidermal differentiation (By similarity). Acts as a negative regulator NF-kappa-B activation through deubiquitination of 'Lys-48'-linked ubiquitination of NFKBIA (PubMed:36651806).

Cellular Location


Nucleus

USP39 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

USP39 Antibody - Images



Western blot analysis of USP39 expression in 293T cell lysate.